

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 June 2001 (21.06.2001)

PCT

(10) International Publication Number
WO 01/44170 A1

(51) International Patent Classification⁷: **C07C 235/46**,
237/30, 233/65, 233/01, 233/88, A61K 31/166, 31/167,
A61P 37/00

(21) International Application Number: PCT/SE00/02505

(22) International Filing Date:
12 December 2000 (12.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9904651-8 17 December 1999 (17.12.1999) SE
0015744.6 27 June 2000 (27.06.2000) GB
0017942.4 22 July 2000 (22.07.2000) GB

(71) Applicant (for all designated States except US): **ASTRAZENECA AB** [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ALCARAZ, Lilian** [FR/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, LE11 5RH (GB). **CAFFREY, Moya** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, LE11 5RH (GB). **FURBER, Mark** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, LE11 5RH (GB). **LUKER, Timothy** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, LE11 5RH (GB). **MORTIMORE, Michael** [GB/GB]; Vertex Pharmaceuticals Ltd., 88

Milton Park Abingdon, Oxfordshire OX14 4RY (GB). **PIMM, Austen** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, LE11 5RH (GB). **THORNE, Phillip** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, LE11 5RH (GB). **WILLIS, Paul** [GB/GB]; AstraZeneca R & D Charnwood, Bakewell Road, Loughborough, LE11 5RH (GB).

(74) Agent: **GLOBAL INTELLECTUAL PROPERTY**; AstraZeneca AB, S-151 85 Södertälje (SE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ADAMANTANE DERIVATIVES



(I)



(II)

(57) Abstract: The invention provides adamantane derivatives of formula (I), a process for their preparation, pharmaceutical compositions containing them, a process for preparing the pharmaceutical compositions, and their use in therapy. In formula (I) D represents CH₂ or CH₂CH₂, E represents C(O)NH or NHC(O) and R³ represents a group of formula (I).


ADAMANTANE DERIVATIVES

The present invention relates to adamantane derivatives, a process for their preparation, pharmaceutical compositions containing them, a process for preparing the pharmaceutical compositions, and their use in therapy.

The P2X₇ receptor (previously known as P2Z receptor), which is a ligand-gated ion channel, is present on a variety of cell types, largely those known to be involved in the inflammatory/immune process, specifically, macrophages, mast cells and lymphocytes (T and B). Activation of the P2X₇ receptor by extracellular nucleotides, in particular adenosine triphosphate, leads to the release of interleukin-1 β (IL-1 β) and giant cell formation (macrophages/microglial cells), degranulation (mast cells) and L-selectin shedding (lymphocytes). P2X₇ receptors are also located on antigen-presenting cells (APC), keratinocytes, salivary acinar cells (parotid cells), hepatocytes, erythrocytes, erythroleukaemic cells, monocytes, fibroblasts, bone marrow cells, neurones and renal mesangial cells.

It would be desirable to make compounds effective as P2X₇ receptor antagonists for use in the treatment of inflammatory, immune or cardiovascular diseases, in the aetiologies of which the P2X₇ receptor may play a role.

In accordance with the present invention, there is therefore provided a compound of general formula




wherein D represents CH₂ or CH₂CH₂, preferably CH₂;

E represents C(O)NH or, preferably, NHC(O);

R¹ and R² each independently represent hydrogen, halogen (e.g. fluorine, chlorine, bromine or iodine), amino (NH₂), nitro (NO₂), C₁-C₆ alkyl or trifluoromethyl, but R¹ and R² may not both simultaneously represent hydrogen;


5 R³ represents a group of formula



R⁴ represents a C₁-C₆ alkyl group;

X represents an oxygen or sulphur atom or a group NR¹³, SO or SO₂;

10 R⁵ represents hydrogen, or R⁵ represents C₁-C₆ alkyl or C₂-C₆ alkenyl, each of which may be optionally substituted by at least one substituent selected from halogen, hydroxyl, (di)-C₁-C₆-alkylamino, -Y-R⁶,



15 a 5- or 6-membered heteroaromatic ring comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulphur which heteroaromatic ring may itself be optionally substituted by at least one substituent selected from halogen, hydroxyl and C₁-C₆ alkyl;

Y represents an oxygen or sulphur atom or a group NH, SO or SO₂;

20 R⁶ represents a group -R⁷Z where R⁷ represents a C₂-C₆ alkyl group and Z represents an -OH, -CO₂H, -NR⁸R⁹, -C(O)NR¹⁰R¹¹ or -N(R¹²)C(O)-C₁-C₆ alkyl group, and,

in the case where Y represents an oxygen or sulphur atom or a group NH, R⁶ additionally represents hydrogen, C₁-C₆ alkyl, C₁-C₆ alkylcarbonyl, C₁-C₆ alkoxycarbonyl, -C(O)NR¹⁴R¹⁵, -CH₂OC(O)R¹⁶, -CH₂OC(O)OR¹⁷ or -C(O)OCH₂OR¹⁸;

25 R⁸, R⁹, R¹⁰, R¹¹ and R¹² each independently represent a hydrogen atom or a C₁-C₆ alkyl group;

R¹³ represents hydrogen, C₃-C₈ cycloalkyl, C₃-C₈ cycloalkylmethyl, or R¹³ represents a C₁-C₆ alkyl group optionally substituted by at least one substituent selected from hydroxyl and C₁-C₆ alkoxy; and

R^{14} , R^{15} , R^{16} , R^{17} and R^{18} each independently represent a C_1 - C_6 alkyl group; with the proviso that when E is $C(O)NH$, X is O, NH or $N(C_1-C_6 \text{ alkyl})$, then R^5 is other than a hydrogen atom or an unsubstituted C_1 - C_6 alkyl group; or a pharmaceutically acceptable salt or solvate thereof.

5

In the context of the present specification, unless otherwise indicated, an alkyl substituent or alkyl moiety in a substituent group may be linear or branched. In the present invention, an alkyl group or moiety may contain up to 6 carbon atoms, examples of which include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, n-pentyl and n-hexyl. A C_2 - C_6 alkenyl group may be linear or branched. In a di- C_1 - C_6 -alkylamino group, the alkyl moieties may be the same or different.

10

In one embodiment of the invention, when E represents $C(O)NH$, then X is S, SO or SO_2 .

15

In another embodiment of the invention, when E represents $NHC(O)$, then X is O or NR^{13} .

20

Preferably, R^1 and R^2 each independently represent a hydrogen or halogen atom, or an amino, nitro, C_1 - C_4 alkyl or trifluoromethyl group (but R^1 and R^2 may not both simultaneously represent a hydrogen atom).

25

More preferably, R^1 and R^2 each independently represent a hydrogen, chlorine or bromine atom, or an amino, nitro, C_1 - C_3 alkyl or trifluoromethyl group (but R^1 and R^2 may not both simultaneously represent a hydrogen atom).

Most preferably, R^1 and R^2 each independently represent a hydrogen or chlorine atom (but R^1 and R^2 may not both simultaneously represent a hydrogen atom).


R^4 represents a C_1 - C_6 alkyl group, for example a linear C_1 - C_6 alkyl group such as CH_2 , $(CH_2)_2$, $(CH_2)_3$ or $(CH_2)_4$.

Preferably X represents an oxygen atom or, especially, a group NR^{13} .

5

R^5 represents hydrogen, or R^5 represents C_1 - C_6 alkyl or C_2 - C_6 alkenyl (e.g. ethenyl or $-CH_2CH=CH_2$), each of which may be optionally substituted by at least one substituent, e.g. one, two or three substituents independently selected from halogen (e.g. fluorine, chlorine, bromine or iodine), hydroxyl, (di)- C_1 - C_6 -alkylamino (e.g. methylamino, ethylamino, dimethylamino or diethylamino), $-Y-R^6$,

10



15

a 5- or 6-membered heteroaromatic ring comprising 1, 2, 3 or 4 heteroatoms independently selected from nitrogen, oxygen and sulphur which heteroaromatic ring may itself be optionally substituted by at least one substituent, e.g. one or two substituents independently selected from halogen (e.g. fluorine, chlorine, bromine or iodine), hydroxyl and C_1 - C_6 , preferably C_1 - C_4 , alkyl, e.g., imidazolyl (such as imidazol-1-yl or imidazol-4-yl), 1-methylimidazolyl (such as 1-methylimidazol-4-yl), 2,3,5-triazolyl and 2,3,4,5-tetrazolyl.

20

Preferred compounds are those in which R^5 represents an optionally substituted C_1 - C_6 alkyl group. A preferred optional substituent is $-Y-R^6$.

25

When Y represents SO or SO_2 , R^6 represents a group $-R^7Z$ where R^7 represents a C_2 - C_6 alkyl group and Z represents an $-OH$, $-CO_2H$, $-NR^8R^9$, $-C(O)NR^{10}R^{11}$ or $-N(R^{12})C(O)-C_1-C_6$ alkyl group.

When Y represents an oxygen or sulphur atom or a group NH, R^6 may represent a group $-R^7Z$ as defined above (particularly $-(CH_2)_2OH$ or $-(CH_2)_3OH$),

or R^6 may represent hydrogen, C_1 - C_6 alkyl (e.g. methyl, ethyl, n-propyl, isopropyl or t-butyl), C_1 - C_6 alkylcarbonyl (e.g. methylcarbonyl, ethylcarbonyl, n-propylcarbonyl or t-butylcarbonyl), C_1 - C_6 alkoxy carbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, n-propoxycarbonyl or t-butoxycarbonyl), $-C(O)NR^{14}R^{15}$, $-CH_2OC(O)R^{16}$,
 5 $-CH_2OC(O)OR^{17}$ or $-C(O)OCH_2OR^{18}$.

Y is preferably an oxygen or sulphur atom or a group NH.

In one embodiment of the invention, Y represents an oxygen or sulphur atom or a
 10 group NH and R^6 represents $-(CH_2)_2OH$, $-(CH_2)_3OH$, hydrogen, methyl, isopropyl, methylcarbonyl or t-butylcarbonyl. In another embodiment, Y represents oxygen and R^6 represents hydrogen.

Preferably R^8 , R^9 , R^{10} , R^{11} and R^{12} each independently represent a hydrogen atom
 15 or a C_1 - C_4 alkyl group.

R^{13} represents hydrogen, C_3 - C_8 , preferably C_3 - C_6 , cycloalkyl, C_3 - C_8 , preferably C_3 - C_6 , cycloalkylmethyl, or R^{13} represents a C_1 - C_6 alkyl group optionally substituted by
 20 at least one substituent, e.g. one, two or three substituents independently selected from hydroxyl and C_1 - C_6 alkoxy. Examples of preferred groups R^{13} include hydrogen, $-(CH_2)_2OH$, methyl, ethyl, n-propyl, isopropyl, n-butyl, n-pentyl, n-hexyl, cyclopropyl, cyclopentyl, cyclohexyl and cyclohexylmethyl.

R^{14} , R^{15} , R^{16} , R^{17} and R^{18} each independently represent a C_1 - C_6 , or C_1 - C_4 , alkyl group.
 25

Preferred compounds of the invention include:

2-Chloro-5-[[2-(2-hydroxy-ethylamino)-ethylamino]-methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
 2-Chloro-5-[[2-(2-hydroxyethoxy)ethylamino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
 30

2-Chloro-5-[(3-hydroxy-2,2-dimethylpropylamino)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[(5-hydroxypentylamino)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5 2-Chloro-5-[[2-[(2-hydroxyethylthio)ethylamino]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(2-hydroxyethyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

10 2-Chloro-5-[3-[(3-hydroxypropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(methylamino)propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(1-methylethyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

15 5-[3-[(2-Amino-2-methylpropyl)amino]propyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(4-hydroxybutyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

20 2-Chloro-5-[3-[(2-hydroxy-2-methylpropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[2-(methylamino)ethyl]amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt,

(*S*)-2-Chloro-5-[3-[(2-hydroxypropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

25 (*R*)-2-Chloro-5-[3-[(2-hydroxypropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

(*R*)-2-Chloro-5-[3-[(2-hydroxy-1-methylethyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

30 2-Chloro-5-[3-[[2-hydroxy-1-(hydroxymethyl)-1-methylethyl]amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5-[3-[[2-(Acetylamino)ethyl]amino]propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[3-[[2-(diethylamino)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt,

5 2-Chloro-5-[3-[(3-methoxypropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[3-[(3-hydroxy-3-methylbutyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

10 2-Chloro-5-[3-[(2-methoxyethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[[3-(methylamino)propoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl) benzamide,

2-Chloro-5-[[2-[(2-hydroxyethyl)amino]ethoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetic acid salt,

15 2-Chloro-5-[[2-[(3-hydroxypropyl)amino]ethoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetic acid salt,

2-Chloro-5-[[[3-[(1-methylethyl)amino]propyl]amino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

20 5-[[[(3-Aminopropyl)amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[[[2-[(1-methylethyl)amino]ethyl]amino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

25 3-[[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]-carbonyl]phenyl]propyl]amino]propanoic acid, 2,2-dimethylpropyl ester, trifluoroacetic acid salt,

5-(2-Aminoethyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide,

2-Chloro-5-[3-[(2-hydroxyethyl)pentylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

30 2-Chloro-5-[3-(methyl-2-propenylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[2-(dimethylamino)ethyl]methylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5-[3-(Butylethylamino)propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5 2-Chloro-5-[3-(methylpentylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[2-(diethylamino)ethyl]ethylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

10 2-Chloro-5-[3-[(2-hydroxyethyl)methylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(dipropylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(2-hydroxyethyl)(1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

15 5-[3-[Butyl(2-hydroxyethyl)amino]propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(diethylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

20 2-Chloro-5-[3-(dimethylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5-[3-(Butylmethylamino)propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(2-hydroxyethyl)propylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

25 2-Chloro-5-[3-[ethyl(2-hydroxyethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(dibutylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

30 2-Chloro-5-[3-(ethylpropylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[methyl(1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[3-(dimethylamino)propyl]methylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5 2-Chloro-5-[3-[cyclohexyl(2-hydroxyethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(cyclohexylmethylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

10 2-Chloro-5-[3-(cyclohexylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[1-(hydroxymethyl)-2,2-dimethylpropyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(cyclopropylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

15 2-Chloro-5-[3-[[2-(dimethylamino)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(3-hydroxy-2,2-dimethylpropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

20 2-Chloro-5-[3-[(1,1-dimethylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[3-(dimethylamino)propyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(cyclopentylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

25 2-Chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-5-[3-[(1,2,2-trimethylpropyl)amino]propyl]-benzamide,

5-[3-(Butylamino)propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[1-(hydroxymethyl)-2-methylpropyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(1-methylpropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[2-(methylthio)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5 2-Chloro-5-[3-[(2-hydroxy-1,1-dimethylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(cyclohexylmethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

10 2-Chloro-5-[3-(2-propenylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(2-fluoroethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(2-methoxy-1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

15 2-Chloro-5-[3-[3-(methylamino)propoxy]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide, dihydrochloride salt,

5-[[[(1-Aminocyclopropyl)methyl](2-hydroxyethyl)amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

20 5-[[[(2-Hydroxyethyl)[2-(methylamino)ethyl]amino]methyl]-2-methyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[2-(1-methyl-1H-imidazol-4-yl)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide,


2-Chloro-5-[3-[[2-(1H-imidazol-4-yl)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide,

25 2-Chloro-5-[3-[[3-(1H-imidazol-1-yl)propyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide,


and pharmaceutically acceptable salts and solvates thereof.

The present invention further provides a process for the preparation of a compound
30 of formula (I) as defined above which comprises:

(a) when X represents an oxygen or sulphur atom or a group NR^{13} , reacting a compound of general formula



wherein L^1 represents a leaving group (e.g. a halogen atom or trifluoromethanesulphonate group) and D, E, R^1 , R^2 and R^4 are as defined in formula (I), with a compound of general formula




wherein X' represents an oxygen or sulphur atom or a group NR^{13} , and R^5 is as defined in formula (I), optionally in the presence of a suitable silver salt (e.g. silver trifluoromethanesulphonate); or

(b) when X represents SO or SO_2 , reacting a corresponding compound of formula (I) in which X represents a sulphur atom with a suitable oxidising agent; or


(c) when X represents a group NR^{13} , reacting a compound of general formula

12



wherein R^{20} represents a bond or C_1 - C_5 alkyl group and D, E, R^1 and R^2 are as defined in formula (I), with a compound of general formula

5




wherein R^5 and R^{13} are as defined in formula (I), in the presence of a reducing agent (e.g. sodium triacetoxyborohydride);

10 and optionally after (a), (b) or (c) converting the compound of formula (I) obtained to a pharmaceutically acceptable salt or solvate thereof.

The processes of the invention may conveniently be carried out in a solvent, e.g. an organic solvent such as dichloromethane, 1, 2-dichloroethane or tetrahydrofuran, at
 15 a temperature, e.g. in the range from 0 to 200 °C, preferably in the range from 0 to 150 °C. The oxidising agent used in (b) above may, for example, be 3-chloroperoxybenzoic acid or potassium peroxymonosulphate, commercially sold under the trade mark "OXONE".

Compounds of formula (V) in which R^{20} represents a bond may be prepared by
 20 reacting a compound of general formula




(VII)

wherein Hal represents a halogen atom such as bromine and D, E, R¹ and R² are as defined in formula (I), with a base such as t-butyllithium and then with a formylating agent such as dimethylformamide.

5


Compounds of formula (VII) may conveniently be prepared by reacting a compound of general formula



(VIII)

wherein R²⁵ represents NH₂ or CO₂H and D is as defined in formula (I),

10 with a compound of general formula




(IX)

wherein R³⁰ represents CO₂H or NH₂, and R¹, R² and Hal are as defined in formula (VII) above.

15

Compounds of formula (V) in which R²⁰ represents a C₁-C₅ alkyl group may be prepared, for example, by reacting a corresponding compound of formula (V) in which R²⁰ represents a bond with (methoxymethyl)diphenylphosphineoxide in the presence of a base, or, with a compound of general formula



in which n is 0, 1, 2 or 3 and R and R' independently represent C₁-C₆ alkyl groups, followed by hydrogenation.


5

Alternatively, compounds of formula (V) in which R²⁰ represents a C₂-C₅ alkyl group may be prepared by reacting a compound of formula (VII) with an alkenol (e.g. 2-propen-1-ol (allyl alcohol), but-3-enol, pent-4-enol or hex-5-enol) in the presence of a palladium catalyst, optionally followed by a hydrogenation reaction and an oxidation
 10 reaction using, for example, Dess-Martin periodinane reagent (these last two steps are not required when the alkenol is allyl alcohol).

As a further alternative, compounds of formula (V) in which R²⁰ represents a C₂-C₅ alkyl group may be prepared by reacting a compound of formula (VII) with an alkenoate
 15 ester (e.g. methyl acrylate or ethyl acrylate) in the presence of a palladium catalyst such as palladium acetate, followed by reduction of the ester group to a hydroxyl group and then oxidation to the aldehyde with an oxidising agent (e.g. Dess-Martin periodinane reagent).


Compounds of formula (I) in which R⁵ represents a C₂-C₆ alkyl group substituted by
 20 a group -Y-R⁶ where Y represents O, S or NH and R⁶ represents a group -R⁷Z as defined above, may be prepared by reacting a corresponding compound of formula (I) in which R⁵ represents a C₂-C₆ alkyl group substituted by a hydroxyl group with a hydroxyl activating agent (such as methanesulphonyl chloride) in the presence of a base (such as triethylamine), followed by reaction with a compound of formula HO-R⁷Z, HS-R⁷Z or
 25 H₂N-R⁷Z.

Compounds of formula (I) wherein E represents a group NHC(O) may be prepared from a compound of general formula



wherein R^1 , R^2 and R^3 are as defined in formula (I) and X represents O, S or NR^{13} , by
 reaction with adamantylmethanamine or adamantylethanamine, in the presence of a
 5 coupling agent such as 1,1'-carbonyldiimidazole.


Compounds of formula (I) wherein E represents a group $C(O)NH$ may be prepared
 from a compound of general formula



10

wherein R^1 , R^2 and R^3 are as defined in formula (I) and X represents O, S or NR^{13} , by
 reaction with adamantylacetyl chloride or adamantylpropanoyl chloride in the presence of a
 base such as triethylamine.


15 Compounds of formula (XI) can be prepared from a compound of general formula



wherein L^2 represents a leaving group (such as a halogen atom or
 trifluoromethanesulphonate group) and R^1 , R^2 and R^4 are as defined in formula (I), with a

compound of formula (IV) as defined above, optionally in the presence of a silver salt such as silver trifluoromethanesulphonate.

Compounds of formula (XII) can be prepared from a compound of general formula



wherein L³ represents a leaving group (such as a halogen atom or trifluoromethanesulphonate group) and R¹, R² and R⁴ are as defined in formula (I), with a compound of formula (IV) as defined above, optionally in the presence of a silver salt such as silver trifluoromethanesulphonate.

10 It will be appreciated that certain compounds of formula (I) may be converted into further compounds of formula (I). For example, compounds of formula (I) in which -Y-R⁶ represents -OH can be converted to compounds in which Y is O and R⁶ is C₁-C₆ alkoxy carbonyl, by reaction with an acylating agent. Furthermore, compounds of formula (I) in which X represents NR¹³ and R¹³ is other than hydrogen, for example, a cyclohexyl group can be prepared by reacting a compound of formula (I) in which X represents NH with cyclohexanone in the presence of a reducing agent such as sodium triacetoxyborohydride.

20 Compounds of formula (III), (IV), (VI), (VIII), (IX), (X), (XIII) and (XIV) as well as compounds HO-R⁷Z, HS-R⁷Z and H₂N-R⁷Z are either commercially available, are well known in the literature or may be prepared easily using known techniques.

25 It will be appreciated by those skilled in the art that in the processes of the present invention certain functional groups such as hydroxyl, carboxyl, aldehyde, carbonyl or amino groups in the starting reagents or intermediate compounds may need to be protected

by protecting groups. Thus, the preparation of the compounds of formula (I) may involve at a certain stage the removal of one or more protecting groups.

The protection and deprotection of functional groups is described in 'Protective Groups in Organic Chemistry', edited by J.W.F. McOmie, Plenum Press (1973) and 'Protective Groups in Organic Synthesis', 2nd edition, T.W. Greene and P.G.M. Wuts, Wiley-Interscience (1991).

The compounds of formula (I) above may be converted to a pharmaceutically acceptable salt or solvate thereof, preferably an acid addition salt such as a hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, tartrate, citrate, oxalate, methanesulphonate or *p*-toluenesulphonate, or an alkali metal salt such as a sodium or potassium salt.

Certain compounds of formula (I) are capable of existing in stereoisomeric forms. It will be understood that the invention encompasses all geometric and optical isomers of the compounds of formula (I) and mixtures thereof including racemates. Tautomers and mixtures thereof also form an aspect of the present invention.

The compounds of the present invention are advantageous in that they possess pharmacological activity and have utility as modulators of P2X₇ receptor activity. They are therefore indicated as pharmaceuticals for use in the treatment or prevention of rheumatoid arthritis, osteoarthritis, psoriasis, allergic dermatitis, asthma, hyperresponsiveness of the airway, chronic obstructive pulmonary disease (COPD), bronchitis, septic shock, glomerulonephritis, irritable bowel disease, Crohn's disease, ulcerative colitis, atherosclerosis, growth and metastases of malignant cells, myoblastic leukaemia, diabetes, neurodegenerative disease, Alzheimer's disease, meningitis, osteoporosis, burn injury, ischaemic heart disease, stroke, peripheral vascular disease and varicose veins.

Accordingly, the present invention provides a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined for use in therapy.

5 In another aspect, the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for use in therapy.

In the context of the present specification, the term "therapy" also includes
10 "prophylaxis" unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

Prophylaxis is expected to be particularly relevant to the treatment of persons who have suffered a previous episode of, or are otherwise considered to be at increased risk of,
15 the disease or condition in question. Persons at risk of developing a particular disease or condition generally include those having a family history of the disease or condition, or those who have been identified by genetic testing or screening to be particularly susceptible to developing the disease or condition.

20 The invention further provides a method of effecting immunosuppression (e.g. in the treatment of rheumatoid arthritis, irritable bowel disease, atherosclerosis, psoriasis, pulmonary disease, e.g. COPD or bronchitis, or diseases of the central nervous system, e.g. Alzheimer's disease or stroke) which comprises administering a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate
25 thereof, as hereinbefore defined to a patient.

For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disease or condition indicated. For effecting immunosuppression, the daily dosage of
30 the compound of formula (I) will typically be in the range from 0.001 mg/kg to 30 mg/kg.

The compounds of formula (I) and pharmaceutically acceptable salts and solvates thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which the formula (I) compound/salt/solvate (active
5 ingredient) is in association with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will preferably comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.10 to 70 %w, of active ingredient, and, from 1 to 99.95 %w, more preferably from 30 to 99.90 %w, of a pharmaceutically acceptable adjuvant, diluent or carrier, all percentages by weight being
10 based on total composition.

Thus, the present invention also provides a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as
15 hereinbefore defined in association with a pharmaceutically acceptable adjuvant, diluent or carrier.


The invention further provides a process for the preparation of a pharmaceutical composition of the invention which comprises mixing a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined with a
20 pharmaceutically acceptable adjuvant, diluent or carrier.

The pharmaceutical composition of the invention may be administered topically (e.g. to the lung and/or airways or to the skin) in the form of solutions, suspensions, heptafluoroalkane aerosols and dry powder formulations; or systemically, e.g. by oral
25 administration in the form of tablets, capsules, syrups, powders or granules, or by parenteral administration in the form of solutions or suspensions, or by subcutaneous administration or by rectal administration in the form of suppositories or transdermally.

The present invention will now be further explained by reference to the following
30 illustrative examples.

Example 1**2-Chloro-5-[[2-(2-hydroxy-ethylamino)-ethylamino]-methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide dihydrochloride**

5

**a) 5-Bromo-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

To a suspension of 5-bromo-2-chlorobenzoic acid (5.00 g) in dichloromethane (25 ml) at 0°C was added oxalyl chloride (3.7 ml) and DMF (5 drops). The resulting mixture was stirred at room temperature under a nitrogen atmosphere for 1h, then concentrated under reduced pressure to yield a solid. The solid was dissolved in dichloromethane (20 ml) and added dropwise to a solution of 1-adamantanemethylamine (3.36g) and *N,N*-diisopropylethylamine (5.55 ml) in dichloromethane (20 ml). The resulting solution was allowed to stir at room temperature under a nitrogen atmosphere for 20hs. The reaction mixture was diluted with dichloromethane and washed with water, 10% aqueous potassium carbonate, 10% aqueous potassium hydrogen sulfate and saturated brine. The organic phase was then dried over sodium sulfate, filtered and concentrated under reduced pressure to afford the subtitled compound as a solid (7.84 g).

20

MS (APCI +ve) 382/384/386 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.42 (1H, t); 7.63 (1H, dd); 7.57 (1H, m); 7.45 (1H, d), 2.93 (2H, d); 1.94 (3H, s, br); 1.69-1.58 (6H, m); 1.51 (6H, s).

b) 2-Chloro-5-formyl-N-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide

5 A solution of 5-bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (3.25 g, Example 1a) in tetrahydrofuran (150 ml) was cooled to -78°C under a nitrogen atmosphere. A solution of 1.4M methyllithium in diethyl ether (6.1 ml) was added to this solution over 2min.. The mixture was stirred at -78°C for 10min., then a 1.7M solution of *tert*-butyllithium in pentane (10.0 ml) was added dropwise. The mixture was stirred at -
10 78°C for a further 10min., then dimethylformamide (1.0 ml) was added. The resulting solution was stirred at -78°C for 30min., quenched with saturated aqueous ammonium chloride solution (100 ml) and extracted with ethyl acetate. The combined extracts were dried over sodium sulfate, filtered, and the filtrate concentrated under reduced pressure to give the subtitled compound as a solid (2.76 g).

15

MS (APCI +ve) 332/334 (M+H)⁺

¹H NMR (DMSO-d₆) δ 10.04 (1H, s); 8.49 (1H, t); 7.96-7.91 (2H, m); 7.74 (1H, d); 2.96 (2H, d), 1.95 (3H, s); 1.64 (6H, m); 1.53 (6H, d).

20 **c) 2-Chloro-5-[[2-(2-hydroxy-ethylamino)-ethylamino]-methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide dihydrochloride**

A mixture of 2-chloro-5-formyl-N-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.244 g, Example 1b), 2-(2-aminoethylamino)-ethanol (0.154 g), p-toluenesulfonic acid (0.005 g) and toluene (30 ml) were refluxed together under Dean-Stark conditions for 3hs,
25 cooled and concentrated under reduced pressure to give an oil. This was dissolved in ethanol (30 ml) and cooled to 0°C under a nitrogen atmosphere. Solid sodium borohydride (0.030 g) was added portionwise to this and the mixture stirred at room temperature for 30min. The mixture was concentrated under reduced pressure and the residue purified by column chromatography over silica gel (eluting with 7:3:0.3 dichloromethane/ methanol/
30 35% aqueous ammonia) to give the free base. This was dissolved in methanol (10 ml) and


treated with 4M hydrochloric acid in dioxane (4 ml) to give a solid precipitate. This was filtered off and washed with diethyl ether to afford the title compound as a solid (0.165 g).

MS (APCI +-ve) 420/422 (M+H)⁺

5 ¹H NMR (DMSO-d₆) δ 8.36 (1H, t); 7.61-7.57 (3H, m); 5.31 (1H s, br); 4.22 (2H, s, br); 3.68 (2H, s, br); 3.05 (2H, s, br); 2.95 (2H, d); 1.95 (3H, s, br); 1.69-1.59 (6H, m); 1.53 (6H, s, br).

Example 2

10 **2-Chloro-5-[[2-(2-hydroxyethoxy)ethylamino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**




15 To a solution of 2-chloro-5-formyl-N-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.150 g, Example 1b) and 2-(2-aminoethoxy)ethanol (0.065 ml) in 1,2-dichloroethane (6 ml) was added sodium triacetoxyborohydride (0.134 g), and the mixture was stirred overnight at room temperature. Water (20 ml) and dichloromethane (20 ml) were added and the layers were partitioned. The organics were washed with brine (30 ml),
20 dried over anhydrous magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by NPHPLC eluting with a gradient of 0-10% ethanol in dichloromethane to give the title compound as a white powder (0.016 g).

MS (APCI +ve) 421/423 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.29 (1H, s, br); 7.42-7.34 (3H, m); 4.60 (1H, s, br); 3.71 (2H, s); 3.47 (4H, s); 3.40 (2H, d); 2.93 (2H, d); 2.63 (2H, d); 1.94 (3H, s); 1.64 (6H, q); 1.52 (6H, s).

5 **Example 3**

2-Chloro-5-[(3-hydroxy-2,2-dimethylpropylamino)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



Prepared according to the method described in Example 2 from 2-chloro-5-formyl-*N*-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.150 g, Example 1b), 3-amino-2,2-dimethylpropanol (0.093 g) and sodium triacetoxyborohydride (0.134 g) in 1,2-dichloroethane (6 ml). After work-up, the residue was purified by NPHPLC eluting with a gradient of 0-10% ethanol in dichloromethane to give the title compound as a white powder (0.035 g).


15

MS (APCI +ve) 419/421 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.29 (1H, t); 7.41-7.34 (3H, m); 4.60 (1H, s, br); 3.70 (2H, s); 3.16 (2H, s); 2.93 (2H, d); 2.29 (2H, s); 1.94 (3H, s, br); 1.63 (6H, q); 1.52 (6H, d); 0.80 (6H, s).

20 **Example 4**

2-Chloro-5-[(5-hydroxypentylamino)methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



A mixture of 2-chloro-5-formyl-*N*-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.100 g, Example 1b), 5-amino-1-pentanol (0.031 mg) and titanium(IV) isopropoxide (0.111 ml) was stirred under nitrogen for 1h at room temperature. The viscous solution obtained was diluted with absolute ethanol (2 ml). Sodium cyanoborohydride (0.013 g) was added, and the solution was stirred for 20h at room temperature. Water (5 ml) was added with stirring and the resulting precipitate was filtered and washed with ethanol. The filtrate was concentrated under reduced pressure. The crude product was dissolved in dichloromethane (20 ml) and filtered to remove the remaining inorganic residues. The filtrate was dried over anhydrous magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by NPHPLC eluting with a gradient of 0-10% ethanol in dichloromethane to give the title compound as a white powder (0.031 g).


MS (APCI +ve) 419/421 (M)⁺

¹H NMR (DMSO-d₆) δ 8.28 (1H, t); 7.41-7.34 (3H, m); 5.30 (1H, s); 4.31 (1H, t); 3.67 (2H, s); 3.37-3.32 (2H, m); 2.93 (2H, d); 2.46-2.42 (2H, m); 1.94 (3H, s); 1.63 (6H, q); 1.52 (6H, s); 1.43-1.36 (2H, m); 1.32-1.28 (2H, m); 1.27-1.21 (2H, m).

Example 5

2-Chloro-5-[[2-(2-hydroxyethylthio)ethylamino]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

25




Prepared according to the method described in Example 2 from 2-chloro-5-formyl-*N*-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.800 g, Example 1b), 2-(2-aminoethylthio)ethanol (0.584 g) and sodium triacetoxyborohydride (0.715 g) in 1,2-dichloroethane (15 ml). After work-up, the residue was purified by NPHPLC eluting with a gradient of 0-10% ethanol in dichloromethane to give the title compound as a white powder (0.536 g).

MS (APCI +ve) 437/439 (M)+

¹H NMR (DMSO-*d*₆) δ 8.29 (1H, t); 7.42-7.35 (3H, m); 4.78 (1H, s, br); 3.71 (2H, s); 3.50 (2H, t); 2.93 (2H, d); 2.68-2.58 (4H, m); 2.56-2.52 (2H, m); 1.94 (3H, s, br); 1.63 (6H, q); 1.52 (6H, s).

Example 6

2-Chloro-5-[3-[(2-hydroxyethyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetate salt



a) (2E)-3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-propenoic acid, methyl ester

5 5-Bromo-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide (5g), methyl acrylate (1.4ml), triethylamine (2.1ml), palladium acetate (0.070g) and tri-orthotolyl phosphine (0.185g) were combined in *N,N*-dimethylformamide (20ml). The mixture was heated in a sealed tube under nitrogen at 90°C for 24h. After cooling, the reaction mixture was partitioned between dichloromethane and dilute hydrochloric acid, the mixed phases were filtered through celite and the phases separated. The organic layer was washed with
10 dilute hydrochloric acid and brine, dried over magnesium sulfate and concentrated under reduced pressure to give a residue which was triturated with diethyl ether and filtered to yield the subtitled compound as an off white solid (4.1g).

MS (APCI +ve) 388/390 (M+H)⁺

15 ¹H NMR (CDCl₃) δ 7.84 (1H, d); 7.64 (1H, d); 7.49 (1H, dd); 7.43 (1H, d); 6.45 (1H, d); 6.24 (1H, t, br); 3.81 (3H, s); 3.19 (2H, d); 2.02 (3H, s); 1.70 (6H, q); 1.59 (6H, d).

b) 4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-benzenepropanoic acid, methyl ester

20 5% Rhodium on carbon (0.40g) was added to a solution of (2E)-3-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-2-propenoic acid, methyl ester (Example 6a, 2.2g) in ethyl acetate / dichloromethane (4:1) (160ml) and the mixture hydrogenated at 3 bar for 24h. The catalyst was removed by filtration and the filtrate concentrated to give the subtitled compound as an oil (2.3g).

25

MS (APCI +ve) 390/392 (M+H)⁺

¹H NMR (CDCl₃) δ 7.55 (1H, d); 7.31 (1H, d); 7.20 (1H, dd); 6.26 (1H, t, br); 3.68 (3H, s); 3.17 (2H, d); 2.95 (2H, t); 2.63 (2H, t); 2.02 (3H, s); 1.70 (6H, q); 1.59 (6H, d).

c) 4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-benzenepropanoic acid

A solution of sodium hydroxide (0.475g) in water (30ml) was added to a solution of 4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-benzenepropanoic acid, methyl ester (Example 6b, 2.3g) in methanol (30ml). After 5h the reaction mixture was reduced to half volume *in vacuo* and acidified with dilute hydrochloric acid. A white solid precipitated and was collected by filtration and dried *in vacuo* at 50°C, to give the subtitled compound (1.2g).

MS (APCI +ve) 376/378 (M+H)⁺

¹H NMR (DMSO-d₆) δ 11.18 (1H, s); 8.28 (1H, t); 7.37 (1H, d); 7.28 (1H, dd); 7.26 (1H, d); 2.92 (2H, d); 2.82 (2H, t); 2.54 (2H, t); 1.94 (3H, s); 1.63 (6H, q); 1.52 (6H, s).

d) 2-Chloro-5-(3-hydroxypropyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Isobutylchloroformate (0.575ml) and triethylamine (0.63ml) were added to a solution of 4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-benzenepropanoic acid (Example 6c, 1.64g) in tetrahydrofuran (30ml) at 0°C. After 1h the precipitates were removed by filtration and the filtrate added portionwise to a solution of sodium borohydride (0.18g) in water (10ml) at 0°C. After a further 1h the reaction mixture was poured onto dilute hydrochloric acid and extracted with ethyl acetate. The organic phase was extracted twice with dilute hydrochloric acid, twice with saturated sodium hydrogencarbonate solution and once with brine, dried over magnesium sulfate and concentrated under reduced pressure to give a residue. Purification by silica gel chromatography (eluting with dichloromethane / methanol 96:4); yielded the subtitled compound as a solid (1.3g).

MS (APCI +ve) 362/364 (M+H)⁺

¹H NMR (CDCl₃) δ 7.55 (1H, d); 7.31 (1H, d); 7.19 (1H, dd); 6.28 (1H, s, br); 3.66 (2H, t); 3.17 (2H, d); 2.72 (2H, t); 1.92 (3H, s); 1.88 (2H, quin); 1.68 (6H, q); 1.59 (6H, s); 1.28 (1H, t).

e) 2-Chloro-5-[3-[(methanesulfonyl)oxy]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

Methanesulfonyl chloride (1.1ml) and triethylamine (2ml) were added to a solution
5 of 2-chloro-5-(3-hydroxypropyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide
(Example 6d, 2.65g) in dichloromethane at 0°C. After 1h the reaction mixture was diluted
with ethyl acetate and extracted once with water, twice with saturated sodium
hydrogencarbonate solution and once with brine, dried over magnesium sulfate and
concentrated under reduced pressure to give the subtitled compound as an oil which slowly
10 solidified (3.2g).

MS (APCI +ve) 440/442 (M+H)⁺

¹H NMR (CDCl₃) δ 7.55 (1H, d); 7.34 (1H, d); 7.20 (1H, dd); 6.32 (1H, t, br); 4.21 (2H, t);
3.18 (2H, d); 3.01 (3H, s); 2.77 (2H, t); 2.09 (2H, quin); 2.01 (3H, s); 1.69 (6H, q); 1.59
15 (6H, d).

f) 2-Chloro-5-[3-[(2-hydroxyethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetate salt


Ethanolamine (0.07ml) was added to a suspension of 2-chloro-5-[3-
20 [(methanesulfonyl)oxy]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example
6e, 0.170g) in n-butanol (5ml) and heated at 100°C in a sealed tube for 12h. On cooling to
ambient temperature the solution was diluted with ethyl acetate and extracted twice with
saturated aqueous sodium hydrogencarbonate solution and once with brine, dried over
magnesium sulfate and concentrated under reduced pressure. Purification by preparative
25 reverse phase HPLC (eluting with a gradient of acetonitrile in 0.1% aqueous ammonium
acetate /25-95%) gave the title compound as the acetate salt (0.070g).

MS (APCI +ve) 405/407 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.28 (1H, t); 7.36 (1H, d); 7.25 (1H, dd); 7.20 (1H, d); 3.44 (2H, t);
30 2.92 (2H, d); 2.50-2.65 (6H, m); 1.94 (3H, s); 1.87 (3H, s); 1.74-1.61 (8H, m); 1.52 (6H, s).

Example 7

2-Chloro-5-[3-[(3-hydroxypropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




3-Aminopropanol (1ml) was added to a solution of 2-chloro-5-[3-[(methylsulfonyl)oxy]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.270g, Example 6e) in tetrahydrofuran (30ml) and the solution heated at reflux for 12h. On cooling to ambient temperature the reaction mixture was diluted with water and extracted thrice with dichloromethane. The organic extracts were combined, dried over magnesium sulfate and concentrated under reduced pressure. Purification by preparative reverse phase HPLC eluting with a gradient of acetonitrile / 0.1% aqueous ammonium acetate (25-95%), gave the title compound as the acetate salt. Treatment with 4M hydrochloric acid in dioxane gave the title compound as the hydrochloride salt (0.070g).

MS (APCI +ve) 419/421 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.67 (2H, s); 8.31 (1H, t); 7.41(1H, d); 7.30-7.25 (2H, m); 4.74 (1H, t); 3.47 (2H, q); 2.95-2.85 (6H, m); 2.67 (2H, t); 2.00-1.84 (5H, m); 1.76 (2H, quin); 1.63 (6H, q); 1.52 (6H, s).

Example 8

2-Chloro-5-[3-(methylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetate salt



Methylamine (2M tetrahydrofuran, 8ml) was added to 2-chloro-5-[3-
[(methylsulfonyl)oxy]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.250g,
Example 6e) and heated in a sealed tube at 70°C for 18h. On cooling to ambient
5 temperature, the solution was diluted with ethyl acetate and extracted twice with saturated
sodium hydrogencarbonate solution and once with brine, dried over magnesium sulfate and
concentrated under reduced pressure. Purification by preparative reverse phase HPLC
(eluting with a gradient of acetonitrile in 0.1% aqueous ammonium acetate /25-95%) gave
the title compound as the acetate salt (0.140g).

10


MS (APCI +ve) 375/377 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.30 (1H, t); 7.37 (1H, d); 7.24 (1H, dd); 7.21 (1H, d); 2.92 (2H,
d); 2.62 (2H, t); 2.53 (2H, t); 2.30 (3H, s); 1.94 (3H, s); 1.86 (3H, s); 1.57-1.77 (8H, m);
1.52 (6H, s).

15

Example 9

2-Chloro-5-[3-[(1-methylethyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




Isopropylamine (0.5ml) was added to a solution of 2-chloro-5-[3-
[(methanesulfonyl)oxy]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.250g,
Example 6e) in tetrahydrofuran (20ml) and heated at 70°C in a sealed tube for 24h. The
mixture was concentrated under reduced pressure and the residue purified by solid phase
extraction on SCX resin. The title product was isolated as the hydrochloric acid salt
(0.10g).

MS (APCI +ve) 403/405 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.67 (2H, s); 8.31 (1H, t); 7.41 (1H, d); 7.30 (1H, dd); 7.26 (1H,
d); 3.33-3.22 (1H, m); 2.93 (2H, d); 2.87 (2H, s); 2.69 (2H, t); 1.86-1.95 (5H, m); 1.63 (6H,
q); 1.52 (6H, s); 1.22 (6H, d).

Example 10

5-[3-[(2-Amino-2-methylpropyl)amino]propyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt




2-Methyl-1,2-propanediamine (0.12ml) was added to a solution of 2-chloro-5-[3-
 [(methylsulfonyl)oxy]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.250g,
 Example 6e) in tetrahydrofuran (4ml) and heated at 60°C in a sealed tube for 12h. On
 5 cooling to ambient temperature, the mixture was diluted with ethyl acetate and extracted
 twice with saturated sodium hydrogencarbonate solution and once with brine, dried over
 magnesium sulfate and concentrated under reduced pressure. Purification by preparative
 reverse phase HPLC (eluting with a gradient of acetonitrile in 0.1% aqueous ammonium
 acetate /25-95%) gave the title compound as the acetate salt. Treatment with 4M
 10 hydrochloric acid in dioxane gave the title compound as the dihydrochloride salt (0.045g).

MS (APCI +ve) 432/434 (M+H)⁺

¹H NMR (DMSO-d₆) δ 9.40 (2H, m); 8.60 (3H, m); 8.32 (1H, t); 7.41 (1H, d); 7.31 (1H,
 d); 7.27 (1H, s); 3.20 (2H, s); 2.92 (4H, d); 2.71 (2H, t); 2.01 (2H, quin); 1.94 (3H, s); 1.63
 15 (6H, q); 1.52 (6H, s); 1.39 (6H, s).

Example 11

**2-Chloro-5-[3-[(4-hydroxybutyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
 benzamide**




4-Amino-1-butanol (0.11 ml) was added to a solution of 2-chloro-5-[3-
 [(methylsulfonyl)oxy]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.25 g,
 Example 6e) in tetrahydrofuran (4 ml) and heated at 60°C in a sealed tube for 12 h. On
 5 cooling to ambient temperature, the mixture was diluted with ethyl acetate and extracted
 twice with saturated aqueous sodium hydrogencarbonate solution and once with brine,
 dried over magnesium sulfate and concentrated under reduced pressure. Purification by
 preparative reverse phase HPLC (eluting with a gradient of acetonitrile in 0.1% aqueous
 ammonium acetate /25-95%) gave the title compound as the acetate salt. Treatment with
 10 aqueous 2M sodium hydroxide and extraction into ethyl acetate gave the title compound
 (0.065 g).

MS (APCI +ve) 433/435 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.28 (1H, t); 7.35 (1H, d); 7.24 (1H, dd); 7.20 (1H, d); 3.37 (2H,
 15 t); 2.92 (2H, d); 2.63 (2H, t); 2.40-2.60 (4H, m); 1.92 (3H, s); 1.70-1.55 (8H, m); 1.52 (6H,
 s); 1.40-1.45 (4H, m).

Example 12

**2-Chloro-5-[3-[(2-hydroxy-2-methylpropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-
 20 ylmethyl)-benzamide, acetate salt**




2-Hydroxy-2-methyl-1-propylamine [prepared according to Journal American Chemical Society (1941), 63, p1034] (0.25ml) was added to a solution of 2-chloro-5-[3-
 5 [(methylsulfonyl)oxy]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.250g, Example 6 e) in butan-1-ol (8ml) and heated at 100°C in a sealed tube for 24h. On cooling to ambient temperature, the mixture was diluted with ethyl acetate and extracted twice with saturated aqueous sodium hydrogencarbonate solution and once with brine, dried over magnesium sulfate and concentrated under reduced pressure. The residue was purified by solid phase extraction on SCX resin and preparative reverse phase HPLC (eluting with a
 10 gradient of acetonitrile in 0.1% aqueous ammonium acetate /25-95%) to give the title compound as the acetate salt (0.160g).

MS (APCI +ve) 433/435 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.29 (1H, t); 7.36 (1H, d); 7.26 (1H, d); 7.21 (1H, s); 2.92 (2H, d);
 15 2.63 (2H, t); 2.55 (2H, t); 2.40 (2H, s); 1.94 (3H, s); 1.88 (3H, s); 1.80-1.58 (8H, m); 1.52 (6H, s); 1.08 (6H, s).

Example 13

2-Chloro-5-[3-[[2-(methylamino)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt
 20



a) 2-Chloro-5-(3-oxopropyl)-benzoic acid,

2-Chloro-5-iodobenzoic acid (5.0g), tetrabutylammonium chloride (5.0g), sodium hydrogencarbonate (5.3g) and allyl alcohol (1.6ml) were combined in N,N-dimethylformamide (50ml) and PdCl₂ (0.6g) was added under nitrogen. After 24h ethyl acetate and 2M hydrochloric acid were added to the crude reaction mixture and the precipitated Pd filtered off. The organic phase was separated and washed thrice with 2M hydrochloric acid then once with brine and dried over magnesium sulfate, filtered and evaporate. Purification by chromatography on silica (eluting ethyl acetate : acetic acid / 19:1) gave the subtitled product as an oil (2.77g).

MS m/z 212/214

¹H NMR (CDCl₃) δ 9.82 (1H, s); 7.83 (1H, d); 7.42 (1H, d); 7.30 (1H, dd); 2.98 (2H, t); 2.83 (2H, t).

b) 2-Chloro-5-[3-[[[(1,1-dimethylethoxy)carbonyl][2-[[[(1,1-dimethylethoxy)carbonyl]methylamino]ethyl]amino]propyl]-benzoic acid,

Sodium triacetoxyborohydride (310mg) was added to a solution of (2-aminoethyl)methyl-, 1,1-dimethylethyl carbamic acid ester [prepared according to J. Med. Chem (1990), 33(1), 100] (0.156g) and 2-chloro-5-(3-oxopropyl)-benzoic acid (0.21g, Example 13a) in methanol (15ml). After 24h, acetic acid (0.2ml) was added and the mixture evaporated to dryness. The residue was redissolved in dichloromethane (10ml), triethylamine (0.45ml) and (1,1-dimethylethoxy)carbonyl 1,1-dimethylethyl

carbonic acid ester (1g) was added. After 24h the reaction mixture was washed twice with 10% KHSO₄, once with brine and dried over magnesium sulfate, filtered and evaporated. Purification by chromatography on silica (eluting with *iso*-hexane : ethyl acetate : acetic acid / 80:20:1 then ethyl acetate : acetic acid / 100:1) gave the subtitled product as an oil
5 (0.2g).

MS (APCI +ve) 471/473 (M+H)⁺

c) 2-Chloro-5-[3-[[2-(methylamino)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt


10 PyBrOP (0.2g) was added to a solution of 2-chloro-5-[3-[[1,1-dimethylethoxy)carbonyl][2-[[1,1-dimethylethoxy)carbonyl]methylamino]ethyl]-amino]propyl]-benzoic acid (0.2g, Example 13b), adamantanemethylamine (0.1ml) and triethylamine (0.15ml) in *N,N*-dimethylformamide (10ml). After 1h the reaction mixture was diluted with ethyl acetate and washed with water, then washed twice with ammonium
15 chloride solution, twice with saturated sodium hydrogencarbonate solution and once with brine. The organic extracts were dried over magnesium sulfate, filtered, evaporated and purified by chromatography on silica (eluting with dichloromethane : methanol /2-10%). The product was redissolved in dichloromethane : methanol / 1:1 (15ml) and 4M hydrochloric acid in dioxane (2ml) added. The mixture was stirred until deprotection was
20 complete then purified by prep RP-HPLC (acetonitrile/0.1% aqueous trifluoroacetic acid). Conversion to the hydrochloride salt by treatment with 4M hydrochloric acid in 1,4-dioxane/methanol gave the title compound (0.055g).

MS (APCI +ve) 418/420 (M+H)⁺

25 ¹H NMR (DMSO-d₆) δ 9.50-9.00 (4H, m); 7.42 (1H, d); 7.29 (1H, d); 7.27 (1H, s); 3.25 (4H, s, br); 3.05-2.90 (4H, m); 2.70 (2H, t); 2.60 (3H, s); 2.00-1.90 (5H, m); 1.63 (6H, q); 1.52 (6H, s).

Example 14

(S)-2-Chloro-5-[3-[(2-hydroxypropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



a) 2-Chloro-5-iodo-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-iodobenzoic acid (10.0g) was suspended in dichloromethane (160ml) then oxalyl chloride (4.0ml) was added followed by *N,N*-dimethylformamide (40μl). After 24 h the solvent was evaporated to afford a white solid, which was redissolved in dichloromethane (160ml). Triethylamine (14.8ml) was added followed by
 10 adamantylmethylamine (6.9ml) with cooling to maintain a temperature below 30°C. The resulting cloudy mixture was stirred for 1h, then evaporated to give a pale yellow solid. This was stirred in a mixture of ethyl acetate (400ml) and 2M hydrochloric acid (300ml) until the solid dissolved to give 2 clear phases. The (upper) organic phase was separated and washed with 2M aqueous sodium hydroxide solution (300ml), then dried (Na₂SO₄) and
 15 evaporated to a yellow solid. The solid was suspended in *iso*-hexane (100ml), then filtered and washed with more *iso*-hexane (40ml). The resulting off-white solid was dried in a vacuum oven at 40°C (14.0g).

MS (APCI +ve) 430/432 (M+H)⁺

20 ¹H NMR (CDCl₃) δ 8.00 (1H, d); 7.66 (1H, dd); 7.14 (1H, d); 6.17 (1H, s, br); 3.17 (2H, d); 2.01 (3H, s); 1.69 (6H, q); 1.58 (6H, d).

b) 2-Chloro-5-(3-oxopropyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

2-Chloro-5-iodo-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (5.00g, Example 14a), tetrabutylammonium chloride (3.40g) and sodium hydrogencarbonate (2.44g) were charged to a flask. Pd(OAc)₂ (0.0533g), toluene (50ml) and allyl alcohol (1.01ml) were added to afford a pale brown mixture which was heated at 80°C for 5h. The resulting dark brown mixture was cooled to ambient then filtered to remove the solid residues. These were washed with further toluene (2 x 50ml) and the combined toluene extracts then washed with water (100ml), dried over MgSO₄ and concentrated to a light brown solid (3.82g).

MS (APCI +ve) 360/362 (M+H)⁺

¹H NMR (CDCl₃) δ 9.81 (1H, s), 7.56 (1H, s); 7.32 (1H, d); 7.19 (1H, d); 6.28 (1H, s, br); 3.18 (2H, d); 2.96 (2H, t); 2.81 (2H, t); 2.01 (3H, s); 1.70 (6H, q); 1.58 (6H, s).

c) **(S)-2-Chloro-5-[3-[(2-hydroxypropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt**


Sodium triacetoxyborohydride (0.6g) was added to a solution of 2-chloro-5-(3-oxopropyl)-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.5g, Example 14b) and (S)-2-hydroxypropylamine (0.31g) in dichloromethane (5ml). After 24h the crude reaction mixture was purified by flash chromatography (eluting with 5-20% methanol in dichloromethane + 1% ammonia) and the hydrochloride salt precipitated from ether/methanol 19:1, to afford the title compound as a white solid (0.19g).

MS (APCI +ve) 419/421 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.70-8.40 (2H, d, br); 8.30 (1H, t); 7.41 (1H, d); 7.28 (1H, dd); 7.24 (1H, d); 5.32 (1H, d); 3.97-3.90 (1H, m); 3.00-2.85 (5H, m); 2.75 (1H, t); 2.65 (2H, t); 2.00-1.90 (5H, m); 1.64 (6H, q); 1.52 (6H, s); 1.09 (3H, d).

Example 15

(R)-2-Chloro-5-[3-[(2-hydroxypropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt




Prepared according to the method described for Example 14.

5 MS (APCI +ve) 419/421 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.70-8.40 (2H, d, br); 8.30 (1H, t); 7.41 (1H, d); 7.28 (1H, dd); 7.24 (1H, d); 5.32 (1H, d); 3.97-3.90 (1H, m); 3.00-2.85 (5H, m); 2.75 (1H, t); 2.65 (2H, t); 2.00-1.90 (5H, m); 1.64 (6H, q); 1.52 (6H, s); 1.09 (3H, d).

10 **Example 16**

(R)-2-Chloro-5-[3-[(2-hydroxy-1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



15 Prepared according to the method described for Example 14.


MS (APCI +ve) 419/421 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.80-8.40 (2H, d, br); 8.31 (1H, t); 7.42 (1H, d); 7.28 (1H, dd); 7.25 (1H, d); 5.36 (1H, s); 3.65-3.60 (1H, d, br); 3.55-3.45 (1H, m); 3.25-3.15 (1H, m); 2.95-2.85 (4H, m); 2.65 (2H, t); 2.00-1.90 (5H, m); 1.63 (6H, q); 1.52 (6H, s); 1.20 (3H, d).

5

Example 17

2-Chloro-5-[3-[[2-hydroxy-1-(hydroxymethyl)-1-methylethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



10

Prepared according to the method described for Example 14.

MS (APCI +ve) 449/451 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.27 (1H, t); 7.36 (1H, d); 7.24 (1H, dd); 7.21 (1H, d); 4.32 (2H, s); 3.22 (4H, s); 2.92 (2H, d); 2.65 (2H, t); 2.45 (2H, q(on edge of DMSO)); 1.92 (3H, s); 1.70-1.57 (8H, m); 1.52 (6H, s); 0.85 (3H, s).


15

Example 18

5-[3-[[2-(Acetylamino)ethyl]amino]propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt

20

41




Prepared according to the method described for Example 14.

MS (APCI +ve) 446/448 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.90 (2H, s, br); 8.32 (1H, t); 8.22 (1H, t); 7.41 (1H, d); 7.28 (1H, dd); 7.24 (1H, d); 3.33 (2H, q); 2.95-2.85 (6H, m); 2.63 (2H, t); 2.00-1.86 (5H, m); 1.84 (3H, s); 1.63 (6H, q); 1.52 (6H, s).

Example 19

2-Chloro-5-[3-[[2-(diethylamino)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt




Prepared according to the method described for Example 14.

MS (APCI +ve) 460/462 (M+H)⁺

¹H NMR (DMSO-d₆) δ 10.70 (1H, s); 9.48 (2H, s); 8.31 (1H, t); 7.42 (1H, d); 7.30 (1H, dd); 7.27 (1H, d); 3.50-3.30 (2H, m); 3.25-3.10 (4H, m); 3.00-2.90 (4H, d, br); 2.71 (2H, t); 2.00-1.90 (5H, m); 1.63 (6H, q); 1.52 (6H, s); 1.24 (6H, t).

5 **Example 20**

2-Chloro-5-[3-[(3-methoxypropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



10 Prepared according to the method described for Example 14.

MS (APCI +ve) 433/435 (M+H)⁺


¹H NMR (DMSO-d₆) δ 8.77 (2H, s, br); 8.31 (1H, t); 7.41 (1H, d); 7.28 (1H, dd); 7.25 (1H, d); 3.40 (2H, t); 3.23 (3H, s); 3.00-2.75 (6H, m); 2.67 (2H, t); 2.00-1.80 (7H, m); 1.63 (6H, q); 1.52 (6H, s).

15

Example 21

2-Chloro-5-[3-[(3-hydroxy-3-methylbutyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt

43




Prepared according to the method described for Example 14.

MS (APCI +ve) (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.57 (2H, s, br); 8.30 (1H, t); 7.42 (1H, d); 7.29 (1H, dd); 7.25 (1H, d); 4.61 (1H, s); 3.05-2.85 (6H, m); 2.64 (2H, t); 2.00-1.82 (5H, m); 1.73-1.53 (8H, m); 1.52 (6H, s); 1.12 (6H, s).

Example 22

2-Chloro-5-[3-[(2-methoxyethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



Prepared according to the method described for Example 14.

15


mp 245-248 °C

MS (APCI +ve) 419/421 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.79 (2H, s); 8.30 (1H, t); 7.41 (1H, d); 7.27 (1H, dd); 7.25 (1H, d); 3.58 (2H, t); 3.30 (3H, s); 3.09 (2H, s, br); 2.95-2.85 (4H, m); 2.67 (2H, t); 1.95-1.86 (5H, m); 1.63 (6H, q); 1.52 (6H, s).

5 **Example 23**

2-Chloro-5-[[3-(methylamino)propoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide



10 **a) 2-Chloro-5-[[3-[[[(1,1-dimethylethoxy)carbonyl]methylamino]propoxy]methyl]-benzoic acid**

(3-Hydroxypropyl)methyl-carbamic acid, 1,1-dimethylethyl ester (0.272 g) in tetrahydrofuran (5 ml) was cooled to 0°C under nitrogen. Sodium hydride (60% dispersion in oil, 0.110 g) was added. The mixture was allowed to warm to room temperature and stirred for 30 min. 5-(Bromomethyl)-2-chloro-benzoic acid (0.300 g) in tetrahydrofuran (3 ml) was added and the mixture heated at 55°C for 6h. The solution was cooled, poured into saturated aqueous potassium hydrogensulfate solution, extracted into ethyl acetate (x 3), dried over magnesium sulfate, filtered and concentrated under reduced pressure to afford the subtitled compound (0.431 g).

20 MS (ESI +ve) 358 (M+H)⁺ (ESI -ve) 356 (M-H)⁺

b) [3-[[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-methoxy]propyl]methyl-carbamic acid, 1,1-dimethylethyl ester

2-Chloro-5-[[3-[[[(1,1-dimethylethoxy)carbonyl]methylamino]propoxy]methyl]-benzoic acid (0.318 g, Example 23a), carbonyl diimidazole (0.165 g) and dimethylformamide (10 ml) were heated at 50°C under nitrogen for 30 min. The mixture was allowed to cool to room temperature and 1-adamantanemethylamine (0.18 ml) was added. The mixture was stirred at room temperature for 20h then poured into ethyl acetate and washed with saturated aqueous sodium hydrogencarbonate solution followed by brine. The solution was dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue was purified by column chromatography over silica gel (eluting with 2:1 isohexane/ethyl acetate) to afford the subtitled compound (0.236 g).

MS (ESI +ve) 505 (M+H)⁺

c) 2-Chloro-5-[[3-(methylamino)propoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide


[3-[[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-methoxy]propyl]methyl-carbamic acid, 1,1-dimethylethyl ester (0.236 g, Example 23b), 4M hydrogen chloride in 1,4-dioxane (5 ml) and methanol (5 ml) were stirred together under nitrogen for 3h, then poured into 25% aqueous ammonia solution and concentrated under reduced pressure to give the free base. This was purified by column chromatography over silica gel, eluting with 19:1:0.1 / dichloromethane:methanol: ammonia to afford the title compound as an oil (0.106 g).

MS (APCI +ve) 405/407 (M+H)⁺

¹H NMR (CDCl₃) δ 7.64 (1H, s); 7.39-7.33 (2H, m); 6.29 (1H, t, br); 4.49 (2H, s); 3.55 (2H, t); 3.18-3.17 (2H, d); 2.72-2.68 (2H, t); 2.44 (3H, s); 2.01 (3H, s, br); 1.86-1.79 (2H, m); 1.75-1.63 (6H, m); 1.59 (6H, s).

Example 24

2-Chloro-5-[[2-[(2-hydroxyethyl)amino]ethoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetic acid salt



a) 5-(Bromomethyl)-2-chloro-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

To a solution of 2-chloro-5-(bromomethyl)-benzoic acid (1.0 g) in dichloromethane (25ml) at 0°C was added dimethylformamide (0.05ml) followed by oxalyl chloride

(0.52 ml). The reaction was allowed to warm to room temperature and stirred for 30min.

The volatiles were removed under vacuum and the residue dried under high vacuum. The

residue was dissolved in dichloromethane (20 ml) and added to a solution of 2-

adamantanemethylamine hydrochloride salt (0.95g) in dichloromethane (20ml) and

diisopropylethylamine (2 ml) at 0°C. The reaction was allowed to warm to room

temperature and stirred for 2h. The organics were washed with water (20ml) then saturated

aqueous ammonium chloride solution and the organic layer dried over magnesium sulfate

then filtered. The filtrate was concentrated under reduced pressure to a solid. The crude

material was recrystallised from dichloromethane/hexane to afford the subtitled compound

as a white solid (1.3 g).

b) [2-[[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl]amino]carbonyl]phenyl]-methoxy]ethyl](2-hydroxyethyl)-carbamic acid, 1,1-dimethylethyl ester

A mixture of 5-(bromomethyl)-2-chloro-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 24a, 0.300g), bis(2-hydroxyethyl)-carbamic acid, 1,1-dimethylethyl

ester (0.312g) and silver trifluoroacetate (0.336 g) in dichloromethane (30 ml) was stirred

under nitrogen at room temperature for 20h. The solution was then decanted from the

silver salts and the dichloromethane removed under vacuum. The crude material was

purified on silica gel (eluting with ethyl acetate) to afford the subtitled compound as an oil

(0.249g).

MS (ESI +ve) MW 521/523 (M+H)⁺

c) 2-Chloro-5-[[2-[(2-hydroxyethyl)amino]ethoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetic acid salt


[2-[[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-methoxy]ethyl](2-hydroxyethyl)-carbamic acid, 1,1-dimethylethyl ester (0.286 g, Example 23b), 4M hydrogen chloride in 1,4-dioxane (10 ml) and methanol (10 ml) were stirred together under nitrogen for 20h, poured into 25% aqueous ammonia solution and concentrated under reduced pressure to give the free base. This was purified by column chromatography over silica gel, eluting with 19:1:0.1 / dichloromethane:methanol: ammonia and further purified by reverse-phase hplc (75:25 to 5:95 / 0.1% aqueous ammonium acetate:acetonitrile) to afford the title compound as an oil (0.051 g).

MS (APCI +ve) 421/423 (M+H)⁺

¹H NMR (CDCl₃) δ 7.66 (1H, d); 7.38 (1H, d); 7.31 (1H dd); 6.68 (1H, t, br); 4.52 (2H, s); 3.72 (2H, t); 3.63 (2H, t); 3.17 (2H, d); 2.95 (2H, t); 2.85 (2H, t); 2.01 (3H, s, br); 1.91 (2H, s); 1.75-1.63 (6H, m); 1.59 (6H, s).

Example 25

2-Chloro-5-[[2-[(3-hydroxypropyl)amino]ethoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetic acid salt



a) 2-Chloro-5-[(2-hydroxyethoxy)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

A mixture of 5-(bromomethyl)-2-chloro-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 24a, 0.300g), ethylene glycol (0.094g) and silver trifluoroacetate (0.336 g) in dichloromethane (10 ml) was stirred under nitrogen at room temperature for 20h. The solution was then decanted from the silver salts and the dichloromethane removed under vacuum. The crude material was purified on silica (eluting with ethyl acetate) to afford the subtitled compound as an oil (0.228g).

MS (ESI +ve) MW 378/380 (M+H)+

b) 2-Chloro-5-[[2-[(methylsulfonyl)oxy]ethoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

2-Chloro-5-[(2-hydroxyethoxy)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.228 g, Example 25a) and triethylamine (0.21 ml) in dichloromethane (10 ml) were cooled to 5°C under nitrogen and methanesulfonyl chloride (0.1 ml) was added. The mixture was stirred at room temperature for 20h then poured into 2N hydrochloric acid, and extracted into ethyl acetate. The combined extracts were washed with 2N hydrochloric acid, saturated aqueous sodium hydrogencarbonate solution then brine, and dried over magnesium sulfate. The organics were filtered and concentrated under reduced pressure to afford the subtitled compound as a white solid (0.160 g).

¹H NMR (CDCl₃) δ 7.66 (1H, s); 7.41-7.33 (2H, m); 6.32 (1H, t, br); 4.57 (2H, s); 4.40-4.38 (2H, m); 3.77-3.75 (2H, m); 3.17-3.15 (2H, d); 3.04 (3H, s); 2.01 (3H, s, br); 1.75-1.64 (6H, m); 1.59 (6H, s, br).

c) 2-Chloro-5-[[2-[(3-hydroxypropyl)amino]ethoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetic acid salt

2-Chloro-5-[[2-[(methylsulfonyl)oxy]ethoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.160 g, Example 25b), 3-amino-1-propanol (0.27 ml) and n-butanol


(4 ml) were heated together in a sealed tube at 110°C for 24h. The mixture was cooled, poured into 2N sodium hydroxide solution and extracted into ethyl acetate. The extracts were dried over sodium sulfate, filtered and concentrated under reduced pressure. Chromatography over silica gel (eluting with 19:1:0.1 / dichloromethane :methanol :ammonia) and further purification by reverse-phase HPLC (75:25 to 5:95 / 0.1% aqueous ammonium acetate:acetonitrile) afforded the title compound as a white solid (0.081 g).

MS (APCI +ve) 435/437 (M+H)⁺

¹H NMR (CDCl₃) δ 7.75 (1H, s); 7.37 (1H, d); 7.27 (1H, d); 6.83 (1H, t, br); 4.56 (2H, s); 3.90 (4H, s, br); 3.21 (4H, s, br); 3.17 (2H, d); 2.01 (5H, s, br); 1.75-1.63 (6H, m); 1.59 (6H, s).

Example 26

2-Chloro-5-[[[3-[(1-methylethyl)amino]propyl]amino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



a) 5-Bromo-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

To a suspension of 5-bromo-2-chlorobenzoic acid (5.00 g) in dichloromethane (25 ml) at 0°C was added oxalyl chloride (3.7 ml) and DMF (5 drops). The resulting mixture was stirred at room temperature under a nitrogen atmosphere for 1h, then concentrated under reduced pressure to yield a solid. The solid was dissolved in dichloromethane (20 ml) and added dropwise to a solution of 1-adamantanemethylamine

(3.36g) and *N,N*-diisopropylethylamine (5.55 ml) in dichloromethane (20 ml). The resulting solution was allowed to stir at room temperature under a nitrogen atmosphere for 20h. The reaction mixture was diluted with dichloromethane and washed in sequence with water, 10% aqueous potassium carbonate solution, 10% aqueous potassium hydrogen sulfate and saturated brine. The organic phase was then dried over sodium sulfate, filtered and concentrated under reduced pressure to afford the subtitled compound as a solid (7.84 g).

MS (APCI +ve) 383/385 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.42 (1H, t); 7.63 (1H, dd); 7.57 (1H, m); 7.45 (1H, d), 2.93 (2H, d); 1.94 (3H, s, br); 1.69-1.58 (6H, m); 1.51 (6H, s).

b) 2-Chloro-5-formyl-*N*-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide

A solution of 5-bromo-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (3.25 g, Example 25a) in tetrahydrofuran (150 ml) was cooled to -78°C under a nitrogen atmosphere. A solution of 1.4M methyllithium in diethyl ether (6.1 ml) was added to this solution over 2min. The mixture was stirred at -78°C for 10min, then a 1.7M solution of *tert*-butyllithium in pentane (10.0 ml) was added dropwise. The mixture was stirred at -78°C for a further 10 min, then dimethylformamide (1.0 ml) was added. The resulting solution was stirred at -78°C for 30 min, quenched with saturated aqueous ammonium chloride solution (100 ml) and extracted with ethyl acetate. The combined extracts were dried over sodium sulfate, filtered, and the filtrate concentrated under reduced pressure to give the subtitled compound as a solid (2.76 g).

MS (APCI +ve) 332/334 (M+H)⁺

¹H NMR (DMSO-d₆) δ 10.04 (1H, s); 8.49 (1H, t); 7.96-7.91 (2H, m); 7.74 (1H, d); 2.96 (2H, d), 1.95 (3H, s); 1.64 (6H, m); 1.53 (6H, d).

c) 2-Chloro-5-[[[3-[(1-methylethyl)amino]propyl]amino]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide


A mixture of 2-chloro-5-formyl-*N*-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.244 g, Example 25b), *N*-isopropyl-1,3-propanediamine (0.211 g), *p*-toluenesulfonic acid (0.005 g) and toluene (30 ml) were refluxed together under Dean-Stark conditions for 3h. The mixture was cooled and concentrated under reduced pressure to an oil. This was dissolved in ethanol (30 ml) and cooled to 0°C under a nitrogen atmosphere. Solid sodium borohydride (0.040 g) was added portionwise and the mixture stirred at room temperature for 30 min, then concentrated under reduced pressure and the residue purified by column chromatography over silica (eluting with 9:1:0.1 dichloromethane/ methanol/ 35% aqueous ammonia) to give the title compound (0.015 g) and a by-product (see Example 27).

MS (APCI +ve) 432/434 (M+H)⁺

¹H NMR (CDCl₃) δ 7.64 (1H, s); 7.37-7.32 (2H, m); 6.28 (1H s, br); 3.78 (2H, s); 3.18 (2H, d); 2.84-2.74 (1H, m); 2.70-2.64 (4H, m); 2.01 (3H, s, br); 1.76-1.63 (8H, m); 1.59 (6H, s, br); 1.05 (6H, d).

Example 27

5-[[[(3-Aminopropyl)amino]methyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




Formed as a by-product in Example 26c above (0.125 g).

MS (APCI +ve) 390/392 (M+H)⁺

¹H NMR (CDCl₃) δ 7.65 (1H, s); 7.37-7.32 (2H, m); 6.31 (1H, t, br); 3.79 (2H, s); 3.18 (2H, d); 2.77 (2H, t); 2.68 (2H, t); 2.01 (3H, s, br); 1.75-1.61 (8H, m); 1.59 (6H, s, br).

Example 28

5 **2-Chloro-5-[[[2-[(1-methylethyl)amino]ethyl]amino]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**




10 Synthesised as in Example 26 using 2-chloro-5-formyl-*N*-(tricyclo [3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide (0.500 g, Example 25b) and *N*-isopropylethylenediamine (0.186 g) to give the title compound as a solid (0.105 g).

MS (APCI +ve) 418/420 (M+H)⁺

15 ¹H NMR (CDCl₃) δ 7.65 (1H, s); 7.34 (2H, s); 6.26 (1H, t, br); 3.79 (2H, s); 3.18 (2H, d); 2.79-2.69 (5H, m); 2.01 (3H, s, br); 1.75-1.64 (6H, m); 1.59 (6H, s, br); 1.50 (2H, s, br), 1.05 (6H, d).

Example 29

20 **3-[[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-propyl]amino]propanoic acid, 2,2-dimethylpropyl ester, trifluoroacetic acid salt**



a) 2-Chloro-5-iodo-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

2-Chloro-5-iodobenzoic acid (10.0g) was suspended in dichloromethane (160ml) then oxalyl chloride (4.0ml) was added followed by *N,N*-dimethylformamide (40μl). After
 5 24 h the solvent was evaporated to afford a white solid, which was then redissolved in dichloromethane (160ml). Triethylamine (14.8ml) was added, followed by adamantane methylamine (6.9ml) with cooling to keep the temperature below 30°C. The resulting slightly cloudy mixture was stirred for 1h, then solvent was evaporated to give a pale yellow solid. The solid was stirred in a mixture of ethyl acetate (400ml) and 2M
 10 hydrochloric acid (300ml) until the solid dissolved to afford two clear phases. The (upper) organic phase was separated off and washed with 2M sodium hydroxide solution (300ml), then dried (Na₂SO₄) and evaporated to a yellow solid. The solid was suspended in *iso*-hexane (100ml), then filtered and washed with more *iso*-hexane (40ml), and the resulting off-white solid dried *in vacuo* at 40°C to afford the subtitled compound (14.0g).

15 MS (APCI +ve) 430/432 (M+H)⁺

¹H NMR (CDCl₃) δ 8.00 (1H, d); 7.66 (1H, dd); 7.14 (1H, d); 6.17 (1H, s, br); 3.17 (2H, d); 2.01 (3H, s); 1.69 (6H, q); 1.58 (6H, d).

b) 2-Chloro-5-(3-oxopropyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-iodo-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (5.00g Example 29a), tetrabutylammonium chloride (3.40g) and sodium hydrogencarbonate (2.44g) were charged to a flask, followed by Pd(OAc)₂ (53.3mg), toluene (50ml) and allyl alcohol (1.01ml). The pale brown mixture was heated at 80°C for 5h, then cooled to ambient

temperature. The mixture was filtered and the residues washed with additional toluene (50ml+50ml). The combined toluene solutions were washed with water (100ml), then dried over MgSO_4 and evaporated to a light brown solid (3.82g).

5 MS (APCI +ve) 360/362 ($\text{M}+\text{H}$)⁺

¹H NMR (CDCl_3) δ 9.81 (1H, s), 7.56 (1H, s); 7.32 (1H, d); 7.19 (1H, d); 6.28 (1H, s, br); 3.18 (2H, d); 2.96 (2H, t); 2.81 (2H, t); 2.01 (3H, s); 1.70 (6H, q); 1.58 (6H, s).

c) **2-Chloro-5-[3-[(3-hydroxypropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**
10

Sodium triacetoxyborohydride (1.86g) was added to a solution of 2-chloro-5-(3-oxopropyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (1.57g, Example 29b) and 3-amino-1-propanol (0.8 ml) in dichloromethane (150ml). After 24h the crude reaction mixture was purified by chromatography (eluting with 5-20% methanol in dichloromethane
15 + 1% ammonia) to give the subtitled compound as a white solid (1.05g).

MS (APCI +ve) 419/421 ($\text{M}+\text{H}$)⁺

d) **3-[[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-phenyl]propyl][(1,1-dimethylethoxy)carbonyl]amino]propanoic acid, 2,2-dimethylpropyl ester**
20

2-Chloro-5-[3-[(3-hydroxypropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.271 g, Example 29c), *N,N*-diisopropylethylamine (0.50 ml) and tetrahydrofuran (10 ml) were stirred together under nitrogen. Trimethylacetyl chloride
25 (0.26 ml) was added dropwise and the mixture stirred at room temperature for 48h, then poured into water and extracted into ethyl acetate. The combined extracts were dried over sodium sulfate, filtered, concentrated under reduced pressure and chromatographed over silica (eluting with 4:1 isohexane/ethyl acetate) to give the subtitled compound (0.158 g) as an oil.

MS (ESI+ve) 603/605 (M+H)⁺

e) 3-[[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-phenyl]propyl]amino]propanoic acid, 2,2-dimethylpropyl ester, trifluoroacetic acid salt


3-[[3-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]-phenyl]propyl][(1,1-dimethylethoxy)carbonyl]amino]propanoic acid, 2,2-dimethylpropyl ester (0.158 g, Example 29d), trifluoroacetic acid (2 ml) and dichloromethane (2 ml) were stirred together under nitrogen for 15 min, then concentrated under reduced pressure to give the title compound (0.170 g).

MS (APCI +ve) 503/505 (M+H)⁺

¹H NMR (CDCl₃) δ 7.40 (1H, d); 7.32 (1H, d); 7.14 (1H dd); 4.12 (2H, t); 3.16 (2H, d); 3.09-2.89 (4H, m); 2.64 (2H, t); 2.06-1.92 (7H, m); 1.75-1.63 (6H, m); 1.58 (6H, s); 1.18 (9H, s).

Example 30

5-(2-Aminoethyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide



a) 2-Chloro-5-(cyanomethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide

Potassium cyanide (525mg) and 18-crown-6 (150mg) were added to a solution of 5-(bromomethyl)-2-chloro-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 24a, 1.8g) in ethanol (50ml) and the mixture heated under reflux for 24h. On cooling the

reaction mixture was evaporated to dryness and the residue partitioned between ethyl acetate and saturated sodium hydrogencarbonate solution. The organics were separated, washed with brine, dried over magnesium sulfate, filtered and evaporated. Purification by chromatography on silica (eluting with a gradient of iso-hexane/ethyl acetate /4:1 to 6:4) gave the subtitled product as a pink solid (1.0g).

MS (APCI +ve) 343/345 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.38 (1H, t); 7.52 (1H, d); 7.43-7.38 (2H, m); 4.08 (2H, s); 2.94 (2H, d); 1.92 (3H, s); 1.63 (6H, q); 1.52 (6H, s).

b) [2-[4-Chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-ethyl]carbamic acid, 1,1-dimethylethyl ester

Sodium borohydride (1.6g) was added portionwise to a cooled solution of 2-chloro-5-(cyanomethyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide (Example 30a, 2.1g), (1,1-dimethylethoxy)carbonyl carbonic acid, 1,1-dimethylethyl ester (2.67g) and cobalt(II)chloride (1.6g) in methanol (100ml). After 1.5h the solvent was evaporated and the residue slurried between ethyl acetate and saturated sodium hydrogencarbonate solution and the insoluble cobalt salts filtered. The organic phase was separated and washed with brine, dried over sodium sulfate, filtered and evaporated to give the subtitled product as a foam (2.2g).

MS (APCI +ve) 447/449 (M+H)⁺

¹H NMR (CDCl₃) δ 7.54 (1H, d); 7.34 (1H, d); 7.19 (1H, dd); 6.28 (1H, s, br); 4.54 (1H, s, br); 3.39 (2H, q); 3.18 (2H, d); 2.80 (2H, t); 2.04 (3H, s); 1.70 (6H, q); 1.64 (6H, s); 1.42 (9H, s).

c) 5-(2-Aminoethyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide, hydrochloride salt

Hydrochloric acid (4M in 1,4-dioxane, 3.0ml) was added to a solution of [2-[4-chloro-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]ethyl]carbamic acid,


1,1-dimethylethyl ester (2.2g, Example 30b) in methanol/dichloromethane (1:1) (50ml). After 24h the solvent was evaporated to leave the title compound as a foam (1.85g).

MS (APCI +ve) 347/349 (M+H)⁺

5 ¹H NMR (DMSO-d₆) δ 8.31 (1H, t); 8.15 (3H, s, br); 7.43 (1H, d); 7.31 (2H, m); 3.05-2.98 (2H, m); 2.95-2.85 (4H, m); 1.94 (3H, s); 1.63 (6H, q); 1.52 (6H, s).

Example 31

10 **2-Chloro-5-[3-[(2-hydroxyethyl)pentylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl) benzamide**




Prepared according to the procedure described in Example 6f

MS (APCI +ve) 475/477 (M+H)⁺

Example 32

2-Chloro-5-[3-(methyl-2-propenylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide



5


Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 415/417 (M+H)⁺

10

Example 33

2-Chloro-5-[3-[[2-(dimethylamino)ethyl]methylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide




Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 446/448 (M+H)⁺

5 **Example 34**

5-[3-(Butylethylamino)propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




10

Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 445/447 (M+H)⁺

Example 35

2-Chloro-5-[3-(methylpentylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




5

Prepared according to the procedure described in 6f

MS (APCI +ve) 445/447 (M+H)⁺

Example 36

10 **2-Chloro-5-[3-[[2-(diethylamino)ethyl]ethylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide**




Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 488/490 (M+H)⁺

Example 37

2-Chloro-5-[3-[(2-hydroxyethyl)methylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide




Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 419/421 (M+H)⁺

Example 38

2-Chloro-5-[3-(dipropylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)- benzamide

62




Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 445/447 (M+H)⁺

5

Example 39

2-Chloro-5-[3-[(2-hydroxyethyl)(1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




10

Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 447/449 (M+H)⁺

15 Example 40

5-[3-[Butyl(2-hydroxyethyl)amino]propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




5

Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 461/463(M+H)⁺

10 Example 41

2-Chloro-5-[3-(diethylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




15 Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 417/419(M+H)⁺

Example 42

2-Chloro-5-[3-(dimethylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
5 benzamide



Prepared according to the procedure described in Example 6f.


10

MS (APCI +ve) 389/391(M+H)⁺

Example 43

5-[3-(Butylmethylamino)propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-
15 benzamide

65




Prepared according to the procedure described in Example 6f.

5 MS (APCI +ve) 431/433(M+H)⁺

Example 44

2-Chloro-5-[3-[(2-hydroxyethyl)propylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

10




Prepared according to the procedure described in Example 6f.

15 MS (APCI +ve) 447/449(M+H)⁺

Example 45

2-Chloro-5-[3-[ethyl(2-hydroxyethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



5


Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 433/435(M+H)⁺

10

Example 46

2-Chloro-5-[3-(dibutylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




Prepared according to the procedure described in Example 6f.

15

MS (APCI +ve) 473/475(M+H)⁺

Example 47

2-Chloro-5-[3-(ethylpropylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



5


Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 431/433(M+H)⁺

10

Example 48

2-Chloro-5-[3-[methyl(1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




15

Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 417/419(M+H)⁺

5 **Example 49**

2-Chloro-5-[3-[[3-(dimethylamino)propyl]methylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



10


Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 460/462(M+H)⁺

15 **Example 50**

2-Chloro-5-[3-[cyclohexyl(2-hydroxyethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

69




Prepared according to the procedure described in Example 6f.

5 MS (APCI +ve) 487/489(M+H)⁺

Example 51

2-Chloro-5-[3-(cyclohexylmethylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



10


Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 457/459(M+H)⁺

15

Example 52

2-Chloro-5-[3-(cyclohexylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 461/463(M+H)⁺

5 **Example 54**

2-Chloro-5-[3-(cyclopropylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




10 Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 401/403(M+H)⁺

Example 55

15 **2-Chloro-5-[3-[[2-(dimethylamino)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

72




Prepared according to the procedure described in Example 6f.

5 MS (APCI +ve) 432/434(M+H)⁺

Example 56

2-Chloro-5-[3-[(3-hydroxy-2,2-dimethylpropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

10




Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 447/449(M+H)⁺

Example 57

2-Chloro-5-[3-[(1,1-dimethylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 417/419(M+H)⁺

Example 58

2-Chloro-5-[3-[[3-(dimethylamino)propyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

74




Prepared according to the procedure described in Example 6f.

5 MS (APCI +ve) 446/448(M+H)⁺

Example 59

2-Chloro-5-[3-(cyclopentylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




10

Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 429/431(M+H)⁺

Example 60

2-Chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-5-[3-[(1,2,2-trimethylpropyl)amino]propyl]-benzamide




5

Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 445/447(M+H)⁺

10 **Example 61**

5-[3-(Butylamino)propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}] dec-1-ylmethyl)-benzamide




Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 417/419(M+H)⁺

5 **Example 62**

2-Chloro-5-[3-[[1-(hydroxymethyl)-2-methylpropyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



10


Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 447/449(M+H)⁺

15 **Example 63**

2-Chloro-5-[3-[(1-methylpropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

77




Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 417/419(M+H)⁺

5

Example 64

2-Chloro-5-[3-[[2-(methylthio)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




10

Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 435/437(M+H)⁺

Example 65

2-Chloro-5-[3-[(2-hydroxy-1,1-dimethylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



5


Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 433/435(M+H)⁺

10

Example 66

2-Chloro-5-[3-[(cyclohexylmethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide




15

Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 457/459(M+H)⁺

Example 67

5 **2-Chloro-5-[3-(2-propenylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**




Prepared according to the procedure described in Example 6f.

10 MS (APCI +ve) 401/403 (M+H)⁺

Example 68

2-Chloro-5-[3-[(2-fluoroethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

80




Prepared according to the procedure described in Example 6f.

5 MS (APCI +ve) 407/409 (M+H)⁺

Example 69

2-Chloro-5-[3-[(2-methoxy-1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide

10




Prepared according to the procedure described in Example 6f.

MS (APCI +ve) 433/435 (M+H)⁺

Example 70

2-Chloro-5-[3-[3-(methylamino)propoxy]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide, dihydrochloride salt



5

a) 2-Chloro-5-(3-iodopropyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl) benzamide

2-Chloro-5-(3-hydroxypropyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (5.9 g, Example 6d), iodine (6.2 g, 24.4 mmol), triphenylphosphine (6.4 g, 24.4 mmol) and imidazole (1.66 g, 24.4 mmol) were stirred in dichloromethane (100ml) at room temperature for 1 h. The reaction was quenched with sat. aq. Na₂S₂O₃ and the layers were separated. The aqueous phase was extracted with ethyl acetate. The combined organic phases were dried (Na₂SO₄), concentrated and purified by column chromatography on silica (eluting with EtOAc : isohexane / 1 : 3) to afford the subtitled compound as a pale yellow solid (6.8 g)

10

MS (APCI +ve) 472/474 (M+H)⁺.

15

b) 2-Chloro-5-[3-[3-(methylamino)propoxy]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide, dihydrochloride salt

Silver trifluoroacetate (0.7 g, 3.2 mmol) was added to 1,1-dimethylethyl (3-hydroxypropyl)methyl carbamic acid (1.8 g, 9.5 mmol) [Synth. Commun. (1995), 25(14), 2135-43] in dichloromethane (20 ml). After 10min. 2-chloro-5-(3-iodopropyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl) benzamide (1.0 g, Example 70a) was added and the mixture was stirred for 24 h. The resulting mixture was washed with water and

20


concentrated. Treatment with 4M hydrochloric acid in 1,4-dioxane/methanol, concentration and purification by RPHPLC gave the title compound (26 mg) as a white solid.

MS (APCI +ve) 433/435 (M+H)⁺

5 ¹H NMR (DMSO-d₆) 8.79-8.35 (1H, m), 8.29 (1H, t, J = 6.2 Hz), 7.38 (1H, d, J = 8.1 Hz), 7.31-7.14 (m, 2H), 3.48-3.31 (4H, m), 2.98-2.81 (4H, m), 2.63 (2H, t, J = 7.6 Hz), 2.50 (3H, s), 1.94 (3H, s), 1.87-1.73 (4H, m), 1.64 (6H, m), 1.52 (6H, m)

Example 71

10 **5-[[[(1-Aminocyclopropyl)methyl](2-hydroxyethyl)amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**



15 a) 5-Bromomethyl-2-chloro-benzoic acid

To a stirred solution of 2-chloro-5-methyl-benzoic acid (25g) in chloroform (500ml) at 50°C was added N-bromosuccinimide (27.40g). The flask was purged with nitrogen and azobisisobutyronitrile (0.10g) added in one portion. The solution was heated at reflux for 1h. Further azobisisobutyronitrile (0.10g) was added and the mixture heated a further 3h.
20 The solution was concentrated in vacuo, redissolved in diethyl ether and filtered to remove insoluble succinimide. The ether solution was washed with 2N aqueous hydrochloric acid solution followed by brine then dried over magnesium sulphate. The solution was concentrated to a volume of 150ml then diluted with isohexane. After further partial

concentration crystallization started. The mixture was allowed to stand in an ice-bath for 1h. The resulting crystals were filtered, washed with isohexane and dried in vacuo to give the subtitled compound (17g).

5 **b) 5-Bromomethyl-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

To a stirred solution of 5-bromomethyl-2-chloro-benzoic acid (Example 71a, 12.4g) in dichloromethane (250ml) and dimethylformamide (0.12ml) at 0°C was added oxalyl chloride (8.7ml). The cooling bath was removed and the solution allowed to warm to room temperature. Once gas evolution had ceased the solution was concentrated *in vacuo*. The
10 residue was redissolved in dichloromethane (300ml), cooled to 0°C and treated with diisopropylethylamine (12.4 ml) and adamantylmethylamine (7.54ml). After 15min. at 0°C the solution was poured into diethyl ether (1L) and washed with 1N aqueous hydrochloric acid followed by brine. The organics were dried over magnesium sulphate and concentrated *in vacuo* to give the title compound as a white powder (19g)

15

MS (APCI +ve) 396/398 (M+H)⁺

¹H NMR (DMSO-d₆) δ 8.39 (1H, t); 7.50-7.40 (2H, m); 4.74 (2H, s); 2.92 (2H, d); 2.50 (3H, s); 1.94 (3H, s, br); 1.67 (3H, d); 1.59 (3H, d); 1.52 (6H, s).

20 **c) 2-Chloro-5-[[[(2-hydroxyethyl)amino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**

A mixture 5-(bromomethyl)-2-chloro-N-(2-tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 71b, 0.300 g) and ethanolamine (0.5 ml) in toluene (40 ml) was heated at 100°C for 24h. The mixture was cooled, poured into saturated aqueous sodium
25 hydrogencarbonate solution, extracted into ethyl acetate, dried over magnesium sulfate, filtered and concentrated under reduced pressure to give the subtitled compound as a gum (0.280 g).

MS (APCI +ve) MW 377/379 (M+H)⁺

d) **[1-[[[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-methyl](2-hydroxyethyl)amino]methyl]cyclopropyl]-carbamic acid, 1,1-dimethylethyl ester**

2-Chloro-5-[[2-hydroxyethyl]amino]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.250 g, Example 71c), (1-formylcyclopropyl)-carbamic acid, 1,1-dimethylethyl ester (0.260 g), sodium triacetoxyborohydride (0.600 g) and dichloromethane (50 ml) were stirred together under nitrogen for 24h. The mixture was poured into saturated aqueous sodium hydrogencarbonate solution, extracted into dichloromethane, dried over magnesium sulfate, filtered and concentrated under reduced pressure. The crude material was purified on silica gel (ethyl acetate), to afford the subtitled compound (0.302 g).

MS (APCI +ve) MW 546/548 (M+H)⁺

e) **5-[[[(1-Aminocyclopropyl)methyl](2-hydroxyethyl)amino]methyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide**


[1-[[[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]-methyl](2-hydroxyethyl)amino]methyl]cyclopropyl]-carbamic acid, 1,1-dimethylethyl ester (Example 71d, 0.302 g) was dissolved in methanol (10 ml) and 4N HCl in dioxane (10 ml) was added. The mixture was stirred for 14h at room temperature, then poured into 25% aqueous ammonia solution and concentrated under reduced pressure to give the free base. This was purified by column chromatography over silica (eluting with 19:1:0.1 dichloromethane/methanol/ ammonia) to afford the title compound as an oil (0.230 g).

MS (APCI +ve) MW 446/448 (M+H)⁺

¹H NMR (CDCl₃) δ 7.69 (1H, s); 7.37 (2H, m); 6.38 (1H, t, br); 3.73 (2H, s); 3.64 (2H, t); 3.18 (2H, d); 2.74 (2H, t); 2.49 (2H, s); 2.21 (3H, s, br); 2.01 (3H, s, br); 1.75-1.64 (6H, m); 1.59 (6H, s, br); 0.59 (2H, t); 0.40 (2H, t).

Example 72

5-[[[(2-Hydroxyethyl)[2-(methylamino)ethyl]amino]methyl]-2-methyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide



5

a) [2-[(2-Hydroxyethyl)[[4-methyl-3-[[[(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]methyl]amino]ethyl]methyl-carbamic acid, 1,1-dimethylethyl ester

10 2-Chloro-5-[[[(2-hydroxyethyl)amino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (0.30 g, Example 71c), methyl(2-oxoethyl)-carbamic acid, 1,1-dimethylethyl ester (0.276 g), sodium triacetoxyborohydride (0.720 g) and dichloromethane (50 ml) were stirred together under nitrogen for 24h. The mixture was poured into saturated aqueous sodium hydrogencarbonate solution, extracted into dichloromethane, dried over
15 magnesium sulfate, filtered and concentrated under reduced pressure. The crude material was purified on silica (eluting with 19:1 dichloromethane/methanol) to afford the subtitled compound (0.285 g).

MS (APCI +ve) MW 534/536 (M+H)+

20

b) 5-[[[(2-Hydroxyethyl)[2-(methylamino)ethyl]amino]methyl]-2-methyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide


[2-[(2-Hydroxyethyl)[[4-methyl-3-[[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]carbonyl]phenyl]methyl]amino]ethyl]methyl-carbamic acid, 1,1-dimethylethyl ester (Example 72a, 0.285 g) was dissolved in methanol (10 ml), 4N HCl in dioxane (10 ml) was added and the mixture stirred for 14h at room temperature. The solution was poured into 25% aqueous ammonia solution and concentrated under reduced pressure to give the free base. This was purified by chromatography over silica gel (eluting with 8:2:0.2 / dichloromethane:methanol: ammonia) to afford the title compound as an oil (0.167 g).

MS (APCI +ve) MW 434/436 (M+H)⁺

¹H NMR (CDCl₃) δ 7.65 (1H, s); 7.35 (2H, s); 6.41 (1H, t, br); 3.67 (2H, s); 3.57 (2H, t); 3.17 (2H, d); 2.68-2.61 (6H, m); 2.38 (3H, s); 2.01 (3H, s, br); 1.76-1.63 (6H, m); 1.60 (6H, s, br).

Example 73

2-Chloro-5-[3-[[2-(1-methyl-1*H*-imidazol-4-yl)ethyl]amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide



Synthesized from 2-chloro-5-[3-[(methylsulfonyl)oxy]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 6e, 0.5g) and 3-methylhistamine (0.22g) according to the procedure described in Example 6f to afford the title compound (0.060g).


MS (APCI +ve) 469/471 (M+H)⁺

¹H NMR (DMSO-d₆) δ 9.45-9.30 (2H, m); 9.02 (1H, s); 8.32 (1H, t); 7.55 (1H, s); 7.41 (1H, d); 7.28 (1H, d); 7.26 (1H, d); 3.82 (3H, s); 3.30-3.20 (2H, m); 3.15 (2H, t); 2.95-2.85 (4H, m); 2.70 (2H, t); 2.02-1.93 (5H, m); 1.63 (6H, q); 1.52 (6H, s).

5

Example 74

2-Chloro-5-[3-[[2-(1*H*-imidazol-4-yl)ethyl]amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide



- 10 Synthesized from 2-chloro-5-[3-[(methylsulfonyl)oxy]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 6e, 1mg) and histamine according to the procedure described in Example 6f to afford the title compound.


MS (APCI +ve) 455/457 (M+H)⁺

15

Example 75

2-Chloro-5-[3-[[3-(1*H*-imidazol-1-yl)propyl]amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide

88




Synthesized from 2-chloro-5-[3-[(methylsulfonyl)oxy]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (Example 6e, 1mg) and 1-(3-aminopropyl)imidazole according to the procedure described in Example 6f to afford the title compound.

MS (APCI +ve) 469/471 (M+H)⁺

Example 76

2-Chloro-5-[3-[(3-hydroxypropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt



Sodium triacetoxyborohydride (4.10g) was added to a solution of 2-chloro-5-(3-oxopropyl)-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide (3.46g, Example 14b) and 3-aminopropanol (1.73ml) in dichloromethane (200ml). After 24h the crude reaction mixture was purified by flash chromatography (eluting with 5-20% methanol /

dichloromethane + 1% ammonia) and the hydrochloride salt precipitated from ether/methanol 19:1, to afford the title compound as a white solid (1.60g).

MS (APCI +ve) 419/421 (M+H)⁺

5 ¹H NMR (DMSO-d₆) δ 8.67 (2H, s); 8.31 (1H, t); 7.41(1H, d); 7.30-7.25 (2H, m); 4.74 (1H, t); 3.47 (2H, q); 2.95-2.85 (6H, m); 2.67 (2H, t); 2.00-1.84 (5H, m); 1.76 (2H, quin); 1.63 (6H, q); 1.52 (6H, s).

Pharmacological Analysis


10 Certain compounds such as benzoylbenzoyl adenosine triphosphate (bbATP) are known to be agonists of the P2X₇ receptor, effecting the formation of pores in the plasma membrane (Drug Development Research (1996), 37(3), p.126). Consequently, when the receptor is activated using bbATP in the presence of ethidium bromide (a fluorescent DNA probe), an increase in the fluorescence of intracellular DNA-bound ethidium bromide is
15 observed. The increase in fluorescence can be used as a measure of P2X₇ receptor activation and therefore to quantify the effect of a compound on the P2X₇ receptor.

In this manner, each of the title compounds of the Examples was tested for antagonist activity at the P2X₇ receptor. Thus, the test was performed in 96-well flat bottomed
20 microtitre plates, the wells being filled with 250 µl of test solution comprising 200 µl of a suspension of THP-1 cells (2.5 x 10⁶ cells/ml) containing 10⁻⁴M ethidium bromide, 25 µl of a high potassium buffer solution containing 10⁻⁵M bbATP, and 25 µl of the high potassium buffer solution containing 3 x 10⁻⁵M test compound. The plate was covered with a plastics sheet and incubated at 37 °C for one hour. The plate was then read in a
25 Perkin-Elmer fluorescent plate reader, excitation 520 nm, emission 595 nm, slit widths: Ex 15 nm, Em 20 nm. For the purposes of comparison, bbATP (a P2X₇ receptor agonist) and pyridoxal 5-phosphate (a P2X₇ receptor antagonist) were used separately in the test as controls. From the readings obtained, a pIC₅₀ figure was calculated for each test compound, this figure being the negative logarithm of the concentration of test compound

necessary to reduce the bbATP agonist activity by 50%. Each of the compounds of the Examples demonstrated antagonist activity, having a pIC_{50} figure > 5.0 .

CLAIMS

1. A compound of general formula




5 wherein D represents CH_2 or CH_2CH_2 ;

E represents $\text{C}(\text{O})\text{NH}$ or $\text{NHC}(\text{O})$;

R^1 and R^2 each independently represent hydrogen, halogen, amino, nitro, C_1 - C_6 alkyl or trifluoromethyl, but R^1 and R^2 may not both simultaneously represent hydrogen;

R^3 represents a group of formula

10




R^4 represents a C_1 - C_6 alkyl group;

X represents an oxygen or sulphur atom or a group NR^{13} , SO or SO_2 ;

R^5 represents hydrogen, or R^5 represents C_1 - C_6 alkyl or C_2 - C_6 alkenyl, each of which

15 may be optionally substituted by at least one substituent selected from halogen, hydroxyl, (di)- C_1 - C_6 -alkylamino, $-\text{Y}-\text{R}^6$,



a 5- or 6-membered heteroaromatic ring comprising from 1 to 4 heteroatoms independently selected from nitrogen, oxygen and sulphur which heteroaromatic ring may itself be

20 optionally substituted by at least one substituent selected from halogen, hydroxyl and C_1 - C_6 alkyl;

Y represents an oxygen or sulphur atom or a group NH , SO or SO_2 ;

R^6 represents a group $-R^7Z$ where R^7 represents a C_2-C_6 alkyl group and Z represents an $-OH$, $-CO_2H$, $-NR^8R^9$, $-C(O)NR^{10}R^{11}$ or $-N(R^{12})C(O)-C_1-C_6$ alkyl group, and, in the case where Y represents an oxygen or sulphur atom or a group NH , R^6 additionally represents hydrogen, C_1-C_6 alkyl, C_1-C_6 alkylcarbonyl, C_1-C_6 alkoxy carbonyl, $-C(O)NR^{14}R^{15}$, $-CH_2OC(O)R^{16}$, $-CH_2OC(O)OR^{17}$ or $-C(O)OCH_2OR^{18}$;

R^8 , R^9 , R^{10} , R^{11} and R^{12} each independently represent a hydrogen atom or a C_1-C_6 alkyl group;

R^{13} represents hydrogen, C_3-C_8 cycloalkyl, C_3-C_8 cycloalkylmethyl, or R^{13} represents a C_1-C_6 alkyl group optionally substituted by at least one substituent selected from hydroxyl and C_1-C_6 alkoxy; and

R^{14} , R^{15} , R^{16} , R^{17} and R^{18} each independently represent a C_1-C_6 alkyl group;

with the proviso that when E is $C(O)NH$, X is O , NH or $N(C_1-C_6 \text{ alkyl})$, then R^5 is other than a hydrogen atom or an unsubstituted C_1-C_6 alkyl group;

or a pharmaceutically acceptable salt or solvate thereof.

15

2. A compound according to claim 1, wherein D represents CH_2 .

3. A compound according to claim 1 or claim 2, wherein E represents $NHC(O)$.

20

4. A compound according to any one of claims 1 to 3, wherein R^1 and R^2 each independently represent a hydrogen, chlorine or bromine atom, or an amino, nitro, C_1-C_3 alkyl or trifluoromethyl group.

25

5. A compound according to any one of the preceding claims, wherein X represents an oxygen atom or a group NR^{13} .

6. A compound according to claim 5, wherein R^{13} represents hydrogen, $-(CH_2)_2OH$, methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *n*-pentyl, *n*-hexyl, cyclopropyl, cyclopentyl, cyclohexyl or cyclohexylmethyl.

30

7. A compound according to any one of the preceding claims, wherein R^5 represents a C_1 - C_6 alkyl group substituted by a single substituent $-Y-R^6$.

8. A compound according to any one of the preceding claims, wherein Y represents an
5 oxygen or sulphur atom or a group NH.

9. A compound according to claim 8, wherein R^6 represents $-(CH_2)_2OH$, $-(CH_2)_3OH$, hydrogen, methyl, isopropyl, methylcarbonyl or t-butylcarbonyl.

10. A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof,
according to claim 1 which is selected from:

2-Chloro-5-[[2-(2-hydroxy-ethylamino)-ethylamino]-methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide dihydrochloride,

15 2-Chloro-5-[[2-(2-hydroxyethoxy)ethylamino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[(3-hydroxy-2,2-dimethylpropylamino)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[(5-hydroxypentylamino)methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

20 2-Chloro-5-[[2-[(2-hydroxyethylthio)ethylamino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(2-hydroxyethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetate salt,

25 2-Chloro-5-[3-[(3-hydroxypropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

2-Chloro-5-[3-(methylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetate salt,

2-Chloro-5-[3-[(1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,

- 5-[3-[(2-Amino-2-methylpropyl)amino]propyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt,
- 2-Chloro-5-[3-[(4-hydroxybutyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
- 5 2-Chloro-5-[3-[(2-hydroxy-2-methylpropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetate salt,
- 2-Chloro-5-[3-[[2-(methylamino)ethyl]amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt,
- (*S*)-2-Chloro-5-[3-[(2-hydroxypropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,
- 10 (*R*)-2-Chloro-5-[3-[(2-hydroxypropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,
- (*R*)-2-Chloro-5-[3-[(2-hydroxy-1-methylethyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,
- 15 2-Chloro-5-[3-[[2-hydroxy-1-(hydroxymethyl)-1-methylethyl]amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,
- 5-[3-[[2-(Acetylamino)ethyl]amino]propyl]-2-chloro-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,
- 2-Chloro-5-[3-[[2-(diethylamino)ethyl]amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, dihydrochloride salt,
- 20 2-Chloro-5-[3-[(3-methoxypropyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,
- 2-Chloro-5-[3-[(3-hydroxy-3-methylbutyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,
- 25 2-Chloro-5-[3-[(2-methoxyethyl)amino]propyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, hydrochloride salt,
- 2-Chloro-5-[3-(methylamino)propoxy]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl) benzamide,
- 2-Chloro-5-[2-[(2-hydroxyethyl)amino]ethoxy]methyl]-*N*-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetic acid salt,
- 30

2-Chloro-5-[[2-[(3-hydroxypropyl)amino]ethoxy]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide, acetic acid salt,

2-Chloro-5-[[[3-[(1-methylethyl)amino]propyl]amino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5 5-[[3-(Aminopropyl)amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[[[2-[(1-methylethyl)amino]ethyl]amino]methyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

10 3-[[3-[4-Chloro-3-[[tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)amino]-carbonyl]phenyl]propyl]amino]propanoic acid, 2,2-dimethylpropyl ester, trifluoroacetic acid salt,

5-(2-Aminoethyl)-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide,

2-Chloro-5-[3-[(2-hydroxyethyl)pentylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

15 2-Chloro-5-[3-(methyl-2-propenylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[2-(dimethylamino)ethyl]methylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

20 5-[3-(Butylethylamino)propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(methylpentylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[2-(diethylamino)ethyl]ethylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

25 2-Chloro-5-[3-[(2-hydroxyethyl)methylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(dipropylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

30 2-Chloro-5-[3-[(2-hydroxyethyl)(1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5-[3-[Butyl(2-hydroxyethyl)amino]propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(diethylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5 2-Chloro-5-[3-(dimethylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5-[3-(Butylmethylamino)propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

10 2-Chloro-5-[3-[(2-hydroxyethyl)propylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[ethyl(2-hydroxyethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(dibutylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

15 2-Chloro-5-[3-(ethylpropylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[methyl(1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

20 2-Chloro-5-[3-[[3-(dimethylamino)propyl]methylamino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[cyclohexyl(2-hydroxyethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-(cyclohexylmethylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

25 2-Chloro-5-[3-(cyclohexylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[1-(hydroxymethyl)-2,2-dimethylpropyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

30 2-Chloro-5-[3-(cyclopropylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[2-(dimethylamino)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(3-hydroxy-2,2-dimethylpropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5 2-Chloro-5-[3-[(1,1-dimethylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[3-(dimethylamino)propyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

10 2-Chloro-5-[3-(cyclopentylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-5-[3-[(1,2,2-trimethylpropyl)amino]propyl]-benzamide,

5-[3-(Butylamino)propyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

15 2-Chloro-5-[3-[[1-(hydroxymethyl)-2-methylpropyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(1-methylpropyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[[2-(methylthio)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

20 2-Chloro-5-[3-[(2-hydroxy-1,1-dimethylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(cyclohexylmethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

25 2-Chloro-5-[3-(2-propenylamino)propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(2-fluoroethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[(2-methoxy-1-methylethyl)amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

2-Chloro-5-[3-[3-(methylamino)propoxy]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide, dihydrochloride salt,

5-[[[(1-Aminocyclopropyl)methyl](2-hydroxyethyl)amino]methyl]-2-chloro-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,

5-[[[(2-Hydroxyethyl)[2-(methylamino)ethyl]amino]methyl]-2-methyl-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)-benzamide,


2-Chloro-5-[3-[[2-(1-methyl-1H-imidazol-4-yl)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide,

2-Chloro-5-[3-[[2-(1H-imidazol-4-yl)ethyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide, and


2-Chloro-5-[3-[[3-(1H-imidazol-1-yl)propyl]amino]propyl]-N-(tricyclo[3.3.1.1^{3,7}]dec-1-ylmethyl)benzamide.

11. A process for the preparation of a compound of formula (I) as defined in claim 1 which comprises:

(a) when X represents an oxygen or sulphur atom or a group NR¹³, reacting a compound of general formula




wherein L¹ represents a leaving group and D, E, R¹, R² and R⁴ are as defined in formula (I), with a compound of general formula




wherein X' represents an oxygen or sulphur atom or a group NR^{13} , and R^5 is as defined in formula (I), optionally in the presence of a suitable silver salt; or

(b) when X represents SO or SO_2 , reacting a corresponding compound of formula (I) in which X represents a sulphur atom with a suitable oxidising agent; or

(c) when X represents a group NR^{13} , reacting a compound of general formula



wherein R^{20} represents a bond or C_1 - C_5 alkyl group and D, E, R^1 and R^2 are as defined in formula (I), with a compound of general formula



wherein R^5 and R^{13} are as defined in formula (I), in the presence of a reducing agent;

and optionally after (a), (b) or (c) converting the compound of formula (I) obtained to a pharmaceutically acceptable salt or solvate thereof.

12. A pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 10 in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

13. A process for the preparation of a pharmaceutical composition as claimed in claim 12 which comprises mixing a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as defined in any one of claims 1 to 10 with a pharmaceutically acceptable adjuvant, diluent or carrier.

14. A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 10 for use in therapy.

15. A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 10 for use in the treatment of rheumatoid arthritis.

16. A compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 10 for use in the treatment of chronic obstructive pulmonary disease.

17. Use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 10 in the manufacture of a medicament for use in therapy.

18. Use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 10 in the manufacture of a medicament for use in treating rheumatoid arthritis.

19. Use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 10 in the manufacture of a medicament for use in treating chronic obstructive pulmonary disease.

20. A method of effecting immunosuppression which comprises administering a therapeutically effective amount of a compound of formula (I), or a pharmaceutically

acceptable salt or solvate thereof, as claimed in any one of claims 1 to 10 to a patient in need thereof.

21. A method of treating rheumatoid arthritis which comprises administering a
5 therapeutically effective amount of a compound of formula (I), or a pharmaceutically
acceptable salt or solvate thereof, as claimed in any one of claims 1 to 10 to a patient in
need thereof.

22. A method of treating chronic obstructive pulmonary disease which comprises
10 administering a therapeutically effective amount of a compound of formula (I), or a
pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 10
to a patient in need thereof.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 00/02505

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07C 235/46, C07C 237/30, C07C 233/65, C07C 233/01, C07C 233/88,
A61K 31/166, A61K 31/167, A61P 37/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07C, A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 9929661 A1 (ASTRA PHARMACEUTICALS LTD.), 17 June 1999 (17.06.99), see the claims --	1-22
X	WO 9929660 A1 (ASTRA PHARMACEUTICALS LTD. ET AL), 17 June 1999 (17.06.99), see especially example 23-25, 31 and the claims --	1-22
P, X	WO 0061569 A1 (ASTRAZENECA AB), 19 October 2000 (19.10.00), see example 66a and the claims --	1-22

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier application or patent but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

15 March 2001

Date of mailing of the international search report

29 -03- 2001

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Solveig Gustavsson/ELY

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 00/02505

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p data-bbox="353 442 1417 611">Journal of Medicinal Chemistry, Volume 15, No 11, 1972, V.L. Narayanan, "Adamantyl Analogs of 2-(3-Dimethylaminopropylthio)cinnamanilide" page 1180 - page 1182</p> <div data-bbox="821 656 995 719" style="text-align: center;"> <p>---</p> <p>-----</p> </div>	1-22

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE00/02505

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **20-22**
because they relate to subject matter not required to be searched by this Authority, namely:
See extra sheet.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE00/02505

Claims 20-22 relates to methods of treatment of the human or animal body by surgery or by therapy/diagnostic methods practised on the human or animal body/ Rule. 39.1.(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compound(s)/composition(s).

INTERNATIONAL SEARCH REPORT

Information on patent family members

25/02/01

International application No.

PCT/SE 00/02505

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9929661	A1	17/06/99	AU	1791399 A	28/06/99
				BR	9813390 A	03/10/00
				EP	1036059 A	20/09/00
				NO	20002786 A	31/07/00
				SE	9704544 D	00/00/00

WO	9929660	A1	17/06/99	AU	1791499 A	28/06/99
				BR	9813368 A	03/10/00
				CN	1280560 T	17/01/01
				EP	1036058 A	20/09/00
				NO	20002785 A	01/08/00
				SE	9704545 D	00/00/00

WO	0061569	A1	19/10/00	AU	3994700 A	14/11/00
				GB	0002330 D	00/00/00
				AU	4950499 A	07/02/00
				NO	20010211 D	00/00/00
				SE	9901270 D	00/00/00
